



**Analysis of Fecal Loadings  
Into Bayous Grande, Chico, and Texar  
Pensacola Bay System, FL**

12 September 2006

Prepared for:  
Florida Department of Health  
Escambia County Health Department

by:

Richard A. Snyder, Ph.D.  
Center for Environmental Diagnostics and Bioremediation  
University of West Florida  
11000 University Parkway  
Pensacola, Florida 32514

(850) 474-2060  
<http://uwf.edu/cedb>



## Table of Contents

|   | <u>Page #</u> |
|---|---------------|
| <b>List of Tables</b> .....   | 3             |
| <b>List of Figures</b> .....  | 4             |
| <b>Abstract</b> .....   | 5             |
| <b>Introduction</b> .....   | 6             |
| <b>Materials and Methods</b> .....  | 7             |
| <b>Results</b> .....  | 15            |
| <i>Bayou Grande</i> .....   | 15            |
| <i>Bayou Chico</i> .....  | 34            |
| <i>Bayou Texar</i> .....  | 55            |
| <b>Site-specific Ground water sampling and Infrared (IR) camera assisted sampling</b> ..... | 74            |
| <b>Discussion and Conclusions</b> .....   | 81            |
| <b>References Cited</b> .....   | 82            |
| <b>Acknowledgements</b> .....   | 82            |

## List of Tables

| Table  | Page # |
|--|--------|
| 1. Station codes and coordinates for Bayou Grande sampling. ....                           | 10     |
| 2. Stations for the analysis of fecal and nutrient concentrations in Bayou Chico, FL ..... | 12     |
| 3. Stations for the analysis of fecal and nutrient concentrations in Bayou Texar, FL. .... | 14     |
| 4. Physico-chemical water quality measures from Bayou Grande sampling. ....                | 16     |
| 5. Summary fecal indicator data from Bayou Grande sampling. ....                           | 17     |
| 6. Regression analysis of Bayou Grande station data as a function of rain .....            | 25     |
| 7. Physico-chemical water quality measures from Bayou Chico sampling. ....                 | 35-36  |
| 8. Summary fecal indicator data from Bayou Chico sampling. ....                            | 37     |
| 9. Regression of Bayou Chico station data as a function of rain. ....                      | 46     |
| 10. Physico-chemical water quality measures from Bayou Texar sampling .....                | 56     |
| 11. Summary fecal indicator data from Bayou Texar sampling. ....                           | 57     |
| 12. Regression of Bayou Texar station data as a function of rain. ....                     | 66     |
| 13. <i>Enterococcus</i> counts for sampling of groundwater in Bayou Grande. ....           | 76     |
| 14. Data from IR assisted sampling in Bayou Chico. ....                                    | 79     |

## List of Figures

| Figure   | Page # |
|--|--------|
| 1. Sampling Station locations in Bayou Grande, FL. ....  | 9      |
| 2. Sampling Station locations in Bayou Chico, FL. ....   | 11     |
| 3. Sampling Station locations in Bayou Texar, FL. ....   | 13     |
| 4. Conservative mixing diagram for Total Phosphate in Bayou Grande . ....                      | 18     |
| 5. Conservative mixing diagram for <i>Enterococcus</i> in Bayou Grande. ....                   | 18     |
| 6. Distribution of mean Phosphate values from Bayou Grande Stations. ....                      | 19     |
| 7. Distribution of mean BOD values from Bayou Grande Stations. ....                            | 20     |
| 8. Distribution of the geomean of <i>Enterococcus</i> counts from Bayou Grande Stations. ....  | 21     |
| 9. Distribution of the maximum of <i>Enterococcus</i> counts from Bayou Grande Stations. ....  | 22     |
| 10. Bayou Grande <i>Enterococcus</i> as a function of rainfall ....                            | 23     |
| 11. Station-specific analysis of rainfall dependence in Bayou Grande. ....                     | 26-31  |
| 12. Slope values from station-specific regression analysis for Bayou Grande ....               | 32     |
| 13. Geomeans of <i>Enterococcus</i> at zero rainfall for Bayou Grande ....                     | 33     |
| 14. Conservative mixing diagrams for nitrogen and phosphorous in Bayou Chico. ....             | 34     |
| 15. Mean Nitrate+Nitrite values for Bayou Chico station data. ....                             | 38     |
| 16. Mean Phosphate values for Bayou Chico station data. ....                                   | 39     |
| 17. Mean BOD values for Bayou Chico station data. ....   | 40     |
| 18. Conservative mixing analysis of <i>Enterococcus</i> in Bayou Chico. ....                   | 41     |
| 19. Geomeans of Bayou Chico station data for <i>Enterococcus</i> counts. ....                  | 42     |
| 20. Bayou Chico station data for <i>Enterococcus</i> minimum counts. ....                      | 43     |
| 21. Bayou Chico station data for <i>Enterococcus</i> maximum counts. ....                      | 44     |
| 22. Bayou Chico <i>Enterococcus</i> as a function of rainfall ....                             | 45     |
| 23. <i>Enterococcus</i> dependence on rainfall by station for Bayou Chico. ....                | 47-52  |
| 24. Distribution of slope values for Bayou Chico station rain dependence. ....                 | 53     |
| 25. Distribution of y-intercept values for Bayou Chico Stations. ....                          | 54     |
| 26. Conservative mixing diagrams for Nitrate + Nitrite and Phosphate in Bayou Texar ....       | 55     |
| 27. Mean Nitrate+Nitrite values in samples for Bayou Texar stations. ....                      | 58     |
| 28. Mean Phosphate values in samples from Bayou Texar stations. ....                           | 59     |
| 29. Mean BOD values in samples from Bayou Texar stations ....                                  | 60     |
| 30. Conservative mixing diagrams for <i>Enterococcus</i> in Bayou Texar ....                   | 61     |
| 31. Geomeans for <i>Enterococcus</i> counts at each station in Bayou Texar. ....               | 62     |
| 32. Minimum <i>Enterococcus</i> counts recorded from each station in Bayou Texar. ....         | 63     |
| 33. Maximum <i>Enterococcus</i> counts recorded from each station in Bayou Texar. ....         | 64     |
| 34. <i>Enterococcus</i> counts from Bayou Texar as a function of rainfall ....                 | 65     |
| 35. Regression analysis of Bayou Texar <i>Enterococcus</i> dependence on rainfall by station . | 67-71  |
| 36. Distribution of slope values for Bayou Texar station rain dependence. ....                 | 72     |
| 37. Distribution of y-intercept estimates for bayou Texar. ....                                | 73     |
| 38. In the north arm of Bayou Grande, a ground water stream ....                               | 76     |
| 39. Visible light and infrared images of the inter tidal beach of Bayou Chico ....             | 78     |
| 40. <i>Enterococcus</i> counts relative to Salinity for IR assisted point sampling.....        | 79     |
| 41. Bayou Chico <i>Enterococcus</i> counts at ground water seeps. ....                         | 80     |

## Abstract

Chronic fecal contamination of waterways within the Pensacola Bay, FL system represents both a public health and environmental problem. This report summarizes the findings of a multi-year study to identify sources of loadings of fecal contamination within the urban bayous of Pensacola, FL: Bayou Grande, Bayou Chico, and Bayou Texar. Thirty-one stations were established along the shoreline of Bayou Grande, forty-two in Bayou Chico, and thirty-three in Bayou Texar. Stations were selected to coincide with storm water drainages, perennial streams, and areas of likely groundwater discharge indicated by topography and freshwater wetland plants in salt water areas. Spatially explicit loading to all three systems was apparent. The intensity of this geographic variability (as variance in system-wide data) increased with moderate rainfall (up to 1.6" within the past 48 hours), but higher levels of rainfall, and presumably associated wind-mixing of the systems, resulted in homogenization of the system and loss of both lower and higher count records. Analysis of station-specific data for rainfall effects on contamination indicated some stations with high concentrations at zero rainfall, presumably from groundwater loadings or feral waterfowl, and others with more dependence on rain, presumably as storm water inputs and enhanced groundwater discharge. In Bayou Grande, the residential areas of the northern and western drainages, and not the Naval Air Station along the southern shore, appear to be the major source areas of chronic fecal contamination to the system. GIS plots indicate older residential development using septic tanks in low-lying areas are source areas. In Bayou Chico, concentrations of nitrogen and fecal bacteria decreased along the salinity gradient of the system as a general trend, indicating the three freshwater and residential areas of the bayou as sources to the system. GIS plots of the data clearly indicate the residential areas, as opposed to the industrial and commercial marina areas along the main part of the bayou, as the sources of contamination. Older residential development using septic tanks in low-lying areas are likely sources. In Bayou Texar, nitrogen and fecal bacteria also decreased along the salinity gradient of the system as a general trend, indicating the Carpenters Creek drainage area as the primary groundwater sources to the system, with the main bayou area served by residential sewer being affected mainly by rainfall. Older residential development using septic tanks and older sewer lines in the Carpenters Creek drainage area are likely sources.

## Introduction

Fecal contamination of surface and ground waters is a nationwide environmental and human health problem. As point sources of this contamination are increasingly restricted and improved, attention is beginning to focus on non-point source loadings from storm water runoff and septic tank (onsite sewage treatment and disposal system; OSTDS) effluents. This focus is being driven by a need to address chronic impairment of water resources that have not responded to increased control of point source discharges. This study was undertaken to attempt to localize and define sources of fecal loading into Bayous Grande, Chico and Texar, the urbanized bayous of the Pensacola Bay system. These bayous have a history of fecal contamination and public health closures for recreational use.

Bayou Grande presents a natural experiment insofar as the northern shoreline and drainage area are largely covered by older residential development using septic tanks systems for wastewater disposal, whereas the southern shoreline and drainage are occupied by the Pensacola Naval Air Station (NAS). This provides for a comparison of residential to undeveloped and lightly developed landscape. NAS is serviced by its own wastewater treatment plant with a surface water effluent discharge near the mouth of the bayou. Much of the shoreline of the NAS is occupied by the base golf course and wooded areas. A northern branch of the bayou extends into a residential area serviced by septic tanks south of Gulf Beach Highway.

Bayou Chico has historically supported more industrial activity than the other urban bayous of the Pensacola Bay system. The upper reaches of the drainage area are bifurcated into a western extension and a northern extension. The latter is further bifurcated into east and west branches. The northern extension passes through two constrictions formed by a former railroad trestle and its earth-filled approaches (south constriction) and a bridge carrying Rt. 98, Navy Boulevard (north constriction). Land use within the Bayou Chico drainage basin is a mix of industrial/commercial and residential development. The north and west reaches of the system are mostly covered by older residential development using septic tanks systems for wastewater disposal. The lower and main part of the bayou is dominated by heavy industry (shipbuilding, scrap metal, chemical manufacturing and distribution, petroleum storage and distribution) and commercial and recreational marinas.

Bayou Texar is surrounded by residential development on a sewer system in contrast to the residential areas around bayous Chico and Grande, and thus provides for a comparison of residential sewer service and septic tank use. Past industrial activity in the Palafox corridor has contributed toxic materials via groundwater flow and storm water runoff to the upper reaches of the bayou. Carpenter's Creek feeding in the northern end of the bayou drains areas using septic tanks. Despite the predominance of sewer service, the bayou has chronic fecal contamination problems.

This report presents the results of multiyear sampling to define spatially explicit sources of fecal contamination into these water bodies.

## Materials and Methods

Stations for sampling were identified by visual survey of the shoreline for storm and groundwater drainage pipes, surface water inputs (intermittent and continuous), and likely areas of ground water discharge, as indicated by freshwater wetland vegetation in salt water areas and land contours. Stations were also established in the open waters of the Bayous.

The distribution of the 31 Bayou Grande stations is displayed in Figure 1. Bayou Grande station names associated station coordinates are listed in Table 1. The distribution of the 42 Bayou Chico stations (37 regularly sampled; 5 added late in the study) is displayed in Figure 2. Bayou Chico station names and coordinates are listed in Table 2. The distribution of the 33 Bayou Texar stations is displayed in Figure 3. Station names and coordinates are listed in Table 3. Bayou Grande samples were taken at monthly intervals from 13 December 1999 to 17 October 2001. Bayou Chico samples were taken at monthly intervals from 11 November 2001 to 30 December 2003. Bayou Texar samples were taken from December 1999 to June 2003.

At each station, time of sampling, water temperature, pH, salinity, and dissolved oxygen were recorded by calibrated water quality meter. Water samples were obtained using State of Florida Department of Environmental Protection (DEP) Standard Operating Procedures (SOP), by Escambia County Health Department (ECHD) personnel. Samples were analyzed by standard methods in either the laboratory of Severn Trent Laboratories, Pensacola, FL (some nutrient analyses) or the Wetlands Research Laboratory at the University of West Florida for Biochemical oxygen demand (BOD; EPA Method 405.1, U.S. EPA, 1983), Enterococci (E; EPA Method 1600, U.S. EPA, 1997), Fecal Coliforms (FC; SM9221E, Eaton, et al., 1995), Total nitrate/nitrite (TNO<sub>3</sub>/NO<sub>2</sub> as N; U.S. EPA, 1993), and Total phosphate (TP as P; U.S. EPA 1983). The UWF Wetlands Lab facility is State of Florida certified for environmental analysis (Lab ID: E71176), conforming to the standards of the National Environmental Laboratory Accreditation Conference (NELAC). The laboratory complies with full chain of custody sample storage and handling practices.

Data were reported to the ECHD as analyses of sample lots (by sampling date) were completed. The final dataset is the subject of this report. Microsoft Excel was used for data reduction and analysis. Limits of detection were reported in lieu of zero values. For each station over the time period of sampling, normality of the data was assessed and arithmetic and geomeans were determined for use as a summary dataset, as appropriate. Standard deviations for arithmetic means and coefficients of variation (standard deviation/mean) for geomeans are reported. Correlation analysis was performed to assess any interrelationships between measured parameters.

Regression analysis was used to determine conservative mixing of measured constituents with seawater. Regression was also used for Log *Enterococcus* counts as a function of rainfall in the 48 hours prior to each sampling event for entire bayous and for each station. Station specific regression models of log *Enterococcus* counts as a function of rainfall were used to estimate geomeans of *Enterococcus* contamination at zero rainfall (y-intercept values) and dependence of contamination on rainfall (slope estimates).

Kaleidagraph (Synergy Software, Inc.) was used to generate graphs. Arcview GIS was used to compile geospatial distribution maps for analytical parameters. Statistical models, graphs, and GIS plots are presented for data visualization purposes. No statistical significance is implied unless clearly stated.

Additional sampling was conducted in Bayous Grande and Chico to isolate ground water concentrations of fecal contamination. Sampling in Bayou Grande visually targeted streams and seepage areas in the intertidal zone at low tides. Sampling in Bayou Chico used a similar approach but with the assistance of an infrared imaging to identify groundwater drainages into the bayou by their thermal signatures. A Flir infrared/visible digital video camera was used to image thermal plumes and record visible images of the same fields. The sampling time was coincident with low surface water temperatures in the bayou and extreme low tides to expose the intertidal zone. Point locations displaying warm groundwater signatures as either general seepage areas or defined rivulets crossing the intertidal zone were sampled for *Enterococcus* analysis. These samples were also analyzed by a molecular source tracking method using the *Bacteroides* assay developed by Bernhard and Field (2000), which has proved to be highly specific for human fecal bacteria (Martin *et al.*, in prep). In many cases, “hotspots” were sampled by digging small holes and allowing the ground water to accumulate and overflow prior to sampling. Data on septic tank and drain field placement were obtained from the public records of septic tank inspections at the Escambia County Health Department and from local residents.

## Bayou Grande Station Codes




Figure 1. Sampling Station locations in Bayou Grande, FL. See Table 1 for latitude and longitude coordinates, and station descriptors.

Table 1. Station codes and coordinates for Bayou Grande sampling

| Station Code | Station Name                | Longitude | Latitude |
|--------------|-----------------------------|-----------|----------|
| 1            | Sherman Grove               | -87.32987 | 30.36418 |
| 2            | Kingsport Ave.              | -87.32744 | 30.36700 |
| 3            | Acapulco Camino             | -87.32447 | 30.36866 |
| 4            | Southside tributary         | -87.31635 | 30.36742 |
| 5            | Athens Ave                  | -87.31509 | 30.37168 |
| 6            | Inlet at Athens             | -87.31378 | 30.37486 |
| 7            | Bremen Ave. tributary       | -87.30502 | 30.37504 |
| 8            | Bartow Ave. tributary       | -87.30037 | 30.37686 |
| 9            | Cousineau Rd. storm drain   | -87.29246 | 30.37464 |
| 10           | Southside site 2            | -87.29843 | 30.36704 |
| 11           | Golf Course drain 1         | -87.28503 | 30.36683 |
| 12           | Greve Rd. tributary         | -87.28598 | 30.37553 |
| 13           | Kalash Dr. storm drain      | -87.28457 | 30.37454 |
| 14           | Navy Point bridge (PC15)    | -87.28157 | 30.38184 |
| 15           | Oak Ave.                    | -87.28064 | 30.38389 |
| 16           | Jamaica Avenue              | -87.28355 | 30.38475 |
| 17           | Syrcler sandbar             | -87.28669 | 30.38302 |
| 18           | Syrcler Dr.                 | -87.28936 | 30.38345 |
| 19           | Loop Road                   | -87.34004 | 30.35609 |
| 20           | Barrios Circle              | -87.34481 | 30.36199 |
| 21           | End of bayou                | -87.35697 | 30.35754 |
| 22           | Fairfield/Rentz creek       | -87.30867 | 30.37946 |
| 23           | Bartow Ave. creek           | -87.29996 | 30.38210 |
| 24           | Paulding Ave. creek         | -87.29645 | 30.38272 |
| 25           | Baublits Rd. SE storm drain | -87.28067 | 30.37678 |
| 26           | Baublits Rd. storm drain    | -87.29115 | 30.37475 |
| 27           | Labree Rd. storm drain      | -87.28975 | 30.37409 |
| 28           | Midbayou                    | -87.29849 | 30.37490 |
| 29           | Navy Point park storm drain | -87.28088 | 30.37571 |
| 30           | Palmettos                   | -87.30014 | 30.36671 |
| 31           | Payne Rd. storm drain       | -87.28355 | 30.37380 |

## Bayou Chico Station Codes




Figure 2. Sampling Station locations in Bayou Chico, FL. See Table 2 for latitude and longitude coordinates, and station descriptors.

Table 2. Stations for the analysis of fecal and nutrient loading in Bayou Chico, FL.

| Station Code | Station Name                 | Longitude | Latitude |
|--------------|------------------------------|-----------|----------|
| 1            | PYC Boat Ramp                | -87.24021 | 30.39940 |
| 2            | Bahia Mar Fuel Dock          | -87.24176 | 30.39968 |
| 3            | Pace Storm Drain             | -87.24248 | 30.40003 |
| 4            | Runyan's Seawall             | -87.24489 | 30.40112 |
| 5            | Scrapyard Phragmites         | -87.24990 | 30.40353 |
| 6            | Midbayou (Scrapyard/Island)  | -87.25202 | 30.40380 |
| 7            | Pensacola Shipyard A-10      | -87.25526 | 30.40657 |
| 8            | Pensacola Shipyard end       | -87.25674 | 30.40838 |
| 9            | Tressle Apartments           | -87.25738 | 30.40922 |
| 10           | Vince Whibbs GMC Storm Drain | -87.25793 | 30.41029 |
| 11           | Navy Boulevard Bridge        | -87.25882 | 30.41142 |
| 12           | Church Fragmites             | -87.25861 | 30.41219 |
| 13           | Sawgrass at Tin Boat House   | -87.25862 | 30.41333 |
| 14           | NE Branch Mouth              | -87.25843 | 30.41520 |
| 15           | NE Branch Midway             | -87.25594 | 30.41633 |
| 16           | NE Branch East End           | -87.25255 | 30.41624 |
| 17           | NW Branch Gazebo             | -87.26085 | 30.41398 |
| 18           | NW end                       | -87.26277 | 30.41391 |
| 19           | Rip Rap                      | -87.25983 | 30.41243 |
| 20           | Juncus at Apartments         | -87.25918 | 30.40960 |
| 21           | Channel Marker 17            | -87.25788 | 30.40589 |
| 22           | Rope Fence                   | -87.26279 | 30.40498 |
| 23           | Lakewood Park                | -87.26125 | 30.40463 |
| 24           | West Branch Cattails         | -87.26509 | 30.40379 |
| 25           | West Branch Marsh Point      | -87.26801 | 30.40154 |
| 26           | West end Last Dock           | -87.26875 | 30.40028 |
| 27           | Swampillies at Green Roof    | -87.26294 | 30.40256 |
| 28           | Tire Pole                    | -87.25742 | 30.40308 |
| 29           | Bell Marine Phragmites       | -87.25329 | 30.40107 |
| 30           | Pelican Pole                 | -87.24731 | 30.40139 |
| 31           | Mahogany Landing             | -87.24649 | 30.39951 |
| 32           | Marker 10/ Pilings           | -87.23941 | 30.39861 |
| 33           | Ditch                        | -87.23859 | 30.39936 |
| 34           | S-Street                     | -87.24606 | 30.41514 |
| 35           | Corry Field Road North       | -87.27490 | 30.41179 |
| 36           | Corry Field Road South       | -87.27374 | 30.39895 |
| 37           | Brigadier                    | -87.28441 | 30.39540 |
| 38           | Fairfield                    | -87.30871 | 30.38973 |
| 39           | Q-Street                     | -87.24393 | 30.41415 |
| 40           | New Warrington               | -87.27674 | 30.39811 |
| 41           | Twin Oaks Apartment          | -87.28008 | 30.41110 |
| 42           | Twin Oaks/Prieto             | -87.28128 | 30.41002 |

# Bayou Texar Station Codes




Figure 3. Sampling Station locations in Bayou Texar, FL. See Table 3 for latitude and longitude coordinates, and station descriptors.

Table 3. Stations for the analysis of fecal and nutrient loading in Bayou Texar, FL.

| Station Code | Station Name                   | Longitude | Latitude |
|--------------|--------------------------------|-----------|----------|
| 1            | Cervantes Bridge               | -87.18822 | 30.42641 |
| 2            | Brainerd St. pond              | -87.19123 | 30.42959 |
| 3            | Bayview Park pvc/storm drain   | -87.18920 | 30.43157 |
| 4            | Tree Roots                     | -87.18696 | 30.43316 |
| 5            | Boathouse (point)              | -87.18630 | 30.43386 |
| 6            | Rocks/Gazebo                   | -87.18692 | 30.43480 |
| 7            | Oriental Garden                | -87.18740 | 30.43598 |
| 8            | South Whaley fragmites         | -87.18806 | 30.43691 |
| 9            | Whaley Ditch storm drain       | -87.18974 | 30.43970 |
| 10           | Birnam Woods green SD          | -87.18915 | 30.44268 |
| 11           | Blackshear Ave. SD             | -87.19229 | 30.44652 |
| 12           | Blanford place FW seep         | -87.20189 | 30.45151 |
| 13           | 34th St. storm drain           | -87.20311 | 30.45383 |
| 14           | Six Cement poles-tan house     | -87.19229 | 30.44652 |
| 15           | Carpenter Creek center         | -87.20573 | 30.45882 |
| 16           | Driftwood 4 SD                 | -87.20410 | 30.45639 |
| 17           | Texar Woods SD                 | -87.19778 | 30.45204 |
| 18           | Seville Dr. (2) SD             | -87.19367 | 30.45011 |
| 19           | Banquos Court SD               | -87.18625 | 30.44256 |
| 20           | Bayou Blvd./Perry SD           | -87.18355 | 30.43212 |
| 21           | 12th Ave. bridge               | -87.20894 | 30.46014 |
| 22           | 9th Ave.                       | -87.21429 | 30.47148 |
| 23           | Airport Blvd.                  | -87.22087 | 30.48151 |
| 24           | Born Court                     | -87.23176 | 30.49196 |
| 25           | Boiling Brook                  | -87.24572 | 30.50169 |
| 26           | Sears Warehouse                | -87.24985 | 30.50461 |
| 27           | Interstate 10-Historical Dist. | -87.23848 | 30.50333 |
| 28           | Olive Road                     | -87.24214 | 30.51103 |
| 29           | Walton/Davis                   | -87.22278 | 30.48440 |
| 30           | Brookside Place                | -87.21309 | 30.46293 |
| 31           | Creekside Office               | -87.21623 | 30.47637 |
| 32           | Springhill                     | -87.21799 | 30.47776 |
| 33           | Burgess Road                   | -87.23502 | 30.49439 |

## Results

### *Bayou Grande*

For Bayou Grande, a total of 757 samples were taken over a time period from 13 December 1999 to 17 October 2001. The summary data are presented in Tables 4 and 5. Samples for Nitrate + Nitrite were processed at two separate facilities, with one reporting a higher detection limit (100 µg/L), which was reported if the samples were not above that level.

None of the physico-chemical parameters could be associated specifically with freshwater inflow to the bayou. Conservative mixing diagrams are often used to identify the behavior of substances along the mixing gradient of freshwater and marine water. In a conservative mixing event, there will be a linear decrease or increase of a substance along the gradient from a freshwater or marine source, respectively. Non-conservative behavior, i.e., non-linearities in this type of plot, indicates either sources (data points above a diagonal dilution line) or sinks (data points below a diagonal dilution line) for a substance along the salinity gradient. In this study, none of the parameters measured behaved absolutely conservatively, especially for Bayou Grande. Figure 4 shows this type of plot for Phosphate. The slope of this line is not significantly different from zero, resulting from samples being below the detection limit for analysis, diffuse sources along the salinity gradient, and/or sediment buffering, as is common for estuaries. Figure 5 shows the analysis for Enterococci. This fecal contamination indicator does not show conservative behavior, indicating that their origin is from diffuse sources along the salinity gradient and not specifically from the freshwater origins of the bayou. Using correlation analysis with the entire dataset (with appropriate transformations for normality), none of the parameters measured were predictive of fecal bacteria concentrations within the system. The only factors showing correlation greater than 70% were variations on the fecal indicators themselves and not between fecal indicators and environmental parameters.

The inability of standard statistical methods applied to the entire dataset to determine fecal contamination concentration patterns within the bayou leads to different ways of viewing the data. Geospatial plotting is a powerful tool in visually determining geographic correlations and loading points that are obscured in traditional statistical approaches. Figures 6-9 show the dataset represented in this fashion using Arcview GIS. Figure 6 displays the distribution of mean Phosphate values, and Figure 7 displays the mean Biological Oxygen Demand (BOD) values for all stations in Bayou Grande. Figure 8 shows the geomean data for *Enterococcus* at each station in the bayou. The high geomean areas for this fecal contamination indicator are found in the wetlands and ditch that parallels Gulf Beach Highway, the upper part of the branch north of Navy Point, and the west end of the Bayou. It is instructive to recognize that the high concentration samples were recovered from developed areas along the northern half of the Bayou drainage basin, whereas relatively low counts were recovered from the open Bayou shoreline and along the undeveloped southern shore. The contrast accounts for mobile sources of fecal contamination that would occur throughout the bayou such as wildlife (raccoons, waterfowl, gulls, etc.). Indeed, wildlife densities would be conceivably higher along the NAS shoreline where less disturbance from human activity would occur. Some non-human mobile sources, however, such as domestic/feral waterfowl fed by waterfront homeowners, gulls attracted to anthropogenic food sources, and dog feces in yards may counter if not overwhelm any contributions by wildlife in undeveloped areas.

Table 4. Physico-chemical water quality measures from Bayou Grande sampling.

| Station Name             | Sta. code | Temp  | std  | pH   | std  | Salinity ppt | std  | DO mg/L | std  | BOD mg/L | CV   | NO <sub>3/2</sub> µg/L | CV   | TP mg/L | CV   |
|--------------------------|-----------|-------|------|------|------|--------------|------|---------|------|----------|------|------------------------|------|---------|------|
| Sherman Grove            | 1         | 22.89 | 6.85 | 7.24 | 0.43 | 19.51        | 7.43 | 6.91    | 1.36 | 2.33     | 1.67 | 57.44                  | 0.16 | 0.103   | 0.07 |
| Kingsport Ave.           | 2         | 23.55 | 6.63 | 7.52 | 0.33 | 21.03        | 6.96 | 7.35    | 1.52 | 2.50     |      | 100                    |      | 0.070   |      |
| Acapulco Camino          | 3         | 23.15 | 7.06 | 7.47 | 0.38 | 21.84        | 6.59 | 6.97    | 2.05 | 1.51     | 0.30 | 40.32                  | 0.17 | 0.098   | 0.04 |
| Southside tributary      | 4         | 22.38 | 7.2  | 7.56 | 0.46 | 20.04        | 8.92 | 7.95    | 1.52 |          |      |                        |      |         |      |
| Athens Ave               | 5         | 23.14 | 6.74 | 7.67 | 0.32 | 22.91        | 6.17 | 7.81    | 1.34 | 2.32     |      | 100                    |      | 0.070   |      |
| Inlet at Athens          | 6         | 23.35 | 6.52 | 7.53 | 0.39 | 22.32        | 6.86 | 7.41    | 1.6  | 1.92     |      | 100                    |      | 0.090   |      |
| Bremen Ave. tributary    | 7         | 23.58 | 6.59 | 7.76 | 0.27 | 23.55        | 5.71 | 7.69    | 1.41 | 1.51     |      | 100                    |      | 0.070   |      |
| Bartow Ave. tributary    | 8         | 23.33 | 6.71 | 7.71 | 0.37 | 22.96        | 6.24 | 7.78    | 1.42 | 1.96     |      | 100                    |      | 0.070   |      |
| Cousineau Rd. SD         | 9         | 23.55 | 6.76 | 7.91 | 0.24 | 23.73        | 5.89 | 8.02    | 1.39 | 2.05     |      | 100                    |      | 0.080   |      |
| Southside site 2         | 10        | 23.02 | 6.71 | 7.8  | 0.26 | 22.96        | 5.85 | 7.87    | 1.42 |          |      |                        |      |         |      |
| Golf Course drain 1      | 11        | 22.67 | 7.11 | 7.77 | 0.28 | 22.15        | 6.22 | 8.13    | 1.24 | 1.75     |      | 100                    |      | 0.060   |      |
| Greve Rd. tributary      | 12        | 24.08 | 6.62 | 7.84 | 0.28 | 22.62        | 6.49 | 7.95    | 1.48 |          |      |                        |      |         |      |
| Kalash Dr. storm drain   | 13        | 23.83 | 6.51 | 7.9  | 0.24 | 23.01        | 6.48 | 8.12    | 1.49 |          |      |                        |      |         |      |
| Navy Pt bridge (PC15)    | 14        | 23.63 | 7.2  | 7.89 | 0.28 | 22.88        | 6.32 | 8.29    | 1.3  | 1.94     |      | 100                    |      | 0.080   |      |
| Oak Ave.                 | 15        | 24.03 | 6.77 | 7.79 | 0.24 | 22.71        | 6.28 | 8.07    | 1.43 | 2.20     | 0.63 | 45.52                  | 0.19 | 0.105   | 0.05 |
| Jamaica Avenue           | 16        | 24.1  | 6.58 | 7.78 | 0.25 | 22.52        | 7.25 | 7.92    | 1.45 | 2.24     |      | 100                    |      | 0.070   |      |
| Syrcl sandbar            | 17        | 24.03 | 7    | 7.88 | 0.17 | 23.43        | 6.63 | 8.08    | 1.42 | 2.05     |      | 100                    |      | 0.070   |      |
| Syrcl Dr.                | 18        | 24.22 | 7.14 | 7.57 | 0.4  | 20.01        | 8.87 | 8.07    | 1.5  | 1.54     |      | 100                    |      | 0.070   |      |
| Loop Road                | 19        | 22.54 | 5.49 | 6.24 | 0.92 | 0.02         | 0.05 | 7.94    | 1.06 | 2.95     | 0.42 | 65.45                  | 0.18 | 0.142   | 0.05 |
| Barrios Circle           | 20        | 26.21 | 7.53 | 6.79 | 0.52 | 11.08        | 7.53 | 7.06    | 2.14 |          |      |                        |      |         |      |
| End of bayou             | 21        | 20.45 | 6.07 | 5.85 | 0.98 | 0.11         | 0.31 | 6.67    | 2.47 | 3.20     |      | 100                    |      | 0.070   |      |
| Fairfield/Rentz creek    | 22        | 20.23 | 6.18 | 5.66 | 1.2  | 0.09         | 0.23 | 3.03    | 1.7  | 3.42     | 0.84 | 63.72                  | 0.21 | 0.123   | 0.15 |
| Bartow Ave. creek        | 23        | 19.83 | 5.34 | 6.04 | 0.91 | 0.04         | 0.05 | 4.97    | 1.51 | 2.64     |      | 100                    |      | 0.090   |      |
| Paulding Ave. creek      | 24        | 20    | 5.17 | 5.72 | 1.17 | 0.03         | 0.05 | 4.07    | 1.5  | 1.78     |      | 100                    |      | 0.090   |      |
| Baublits Rd. SE SD       | 25        | 24.11 | 6.88 | 7.88 | 0.28 | 22.52        | 6.63 | 8.34    | 1.25 |          |      |                        |      |         |      |
| Baublits Rd. storm drain | 26        | 23.77 | 6.64 | 7.85 | 0.31 | 23.37        | 6.06 | 7.95    | 1.51 |          |      |                        |      |         |      |
| Labree Rd. storm drain   | 27        | 23.88 | 6.69 | 7.86 | 0.27 | 23.35        | 5.95 | 8       | 1.36 |          |      |                        |      |         |      |
| Midbayou                 | 28        | 23.22 | 6.52 | 7.85 | 0.76 | 23.66        | 12.6 | 7.99    | 2.17 | 1.72     |      | 100                    |      | 0.070   |      |
| Navy Pt park SD          | 29        | 24.08 | 7    | 7.9  | 0.31 | 22.57        | 6.73 | 13.1    | 17.7 |          |      |                        |      |         |      |
| Palmettos                | 30        | 23.4  | 6.73 | 7.81 | 0.24 | 22.2         | 6.32 | 7.73    | 1.58 |          |      |                        |      |         |      |
| Payne Rd. storm drain    | 31        | 23.75 | 6.76 | 7.89 | 0.33 | 23           | 6.44 | 8.32    | 1.41 |          |      |                        |      |         |      |

Table 5. Summary fecal indicator data from Bayou Grande sampling.

| station name                | station code | Entero GeoMean | Entero CV | Entero Max | Entero Min | Fecal Geomean | Fecal CV | Fecal Max | Fecal Min |
|-----------------------------|--------------|----------------|-----------|------------|------------|---------------|----------|-----------|-----------|
| Sherman Grove               | 1            | 42.11          | 0.44      | 2830       | 2          | 68.32         | 0.33     | 4400      | 20        |
| Kingsport Ave.              | 2            | 45.96          | 0.41      | 230        | 4          | 836.52        | 0.29     | 9000      | 20        |
| Acapulco Camino             | 3            | 78             | 0.39      | 6560       | 4          | 471.67        | 0.35     | 9000      | 20        |
| Southside tributary         | 4            | 15.49          | 0.5       | 300        | 1          | 149.36        | 0.33     | 1700      | 20        |
| Athens Ave                  | 5            | 12.53          | 0.67      | 170        | 1          | 139.55        | 0.37     | 1700      | 20        |
| Inlet at Athens             | 6            | 26.82          | 0.52      | 1700       | 1          | 265.45        | 0.36     | 3000      | 20        |
| Bremen Ave. tributary       | 7            | 9.35           | 0.71      | 130        | 1          | 47.27         | 0.23     | 170       | 20        |
| Bartow Ave. tributary       | 8            | 22.2           | 0.57      | 490        | 1          | 187.27        | 0.36     | 1300      | 20        |
| Cousineau Rd. storm drain   | 9            | 6.09           | 0.7       | 56         | 1          | 10.5          |          | 1         | 1         |
| Southside site 2            | 10           | 7              | 0.66      | 52         | 1          | 1.45          | 0.26     | 589       | 20        |
| Golf Course drain 1         | 11           | 19.88          | 0.61      | 500        | 1          | 58.57         | 0.24     | 500       | 20        |
| Greve Rd. tributary         | 12           | 20.64          | 0.47      | 240        | 1          |               |          |           |           |
| Kalash Dr. storm drain      | 13           | 7.5            | 0.94      | 2000       | 1          |               |          |           |           |
| Navy Point bridge (PC15)    | 14           | 10.57          | 0.57      | 230        | 1          | 91.9          | 0.27     | 800       | 20        |
| Oak Ave.                    | 15           | 75.59          | 0.42      | 9000       | 11         | 1198.95       | 0.29     | 16000     | 20        |
| Jamaica Avenue              | 16           | 39.54          | 0.41      | 2210       | 3          | 681.82        | 0.35     | 9000      | 20        |
| Syrclle sandbar             | 17           | 16.6           | 0.57      | 600        | 1          | 811.36        | 0.38     | 16000     | 20        |
| Syrclle Dr.                 | 18           | 67.39          | 0.38      | 3000       | 4          | 482.73        | 0.31     | 3000      | 20        |
| Loop Road                   | 19           | 50.62          | 0.37      | 1300       | 1          | 235.24        | 0.32     | 2400      | 20        |
| Barrios Circle              | 20           | 43.15          | 0.33      | 230        | 4          | 868.64        | 0.29     | 9000      | 20        |
| End of bayou                | 21           | 45.07          | 0.35      | 500        | 4          | 301.5         | 0.36     | 1700      | 20        |
| Fairfield/Rentz creek       | 22           | 115.76         | 0.26      | 600        | 13         | 1984          | 0.39     | 16000     | 20        |
| Bartow Ave. creek           | 23           | 111.86         | 0.31      | 1210       | 19         | 879.41        | 0.36     | 9000      | 20        |
| Paulding Ave. creek         | 24           | 257.09         | 0.21      | 1520       | 20         | 2758.18       | 0.31     | 16000     | 20        |
| Baublits Rd. SE storm drain | 25           | 4.24           | 1.22      | 200        | 1          |               |          |           |           |
| Baublits Rd. storm drain    | 26           | 7.7            | 0.87      | 880        | 1          |               |          |           |           |
| Labree Rd. storm drain      | 27           | 9.04           | 0.86      | 280        | 1          |               |          |           |           |
| Midbayou                    | 28           | 4.3            | 0.97      | 28         | 1          | 28.23         | 0.28     | 130       | 1         |
| Navy Point park storm drain | 29           | 3.8            | 1.14      | 102        | 1          |               |          |           |           |
| Palmettos                   | 30           | 5.29           | 0.83      | 44         | 1          | 56.67         | 0.26     | 500       | 20        |
| Payne Rd. storm drain       | 31           | 6.2            | 1.07      | 2000       | 1          |               |          |           |           |




Figure 4. Conservative mixing diagram for Total Phosphate in Bayou Grande.




Figure 5. Conservative mixing diagram for *Enterococcus* in Bayou Grande.

# Bayou Grande Total Phosphorous (mg/L)




Figure 6. Distribution of mean Phosphate values from Bayou Grande Stations.




Figure 7. Distribution of mean BOD values from Bayou Grande Stations.

Bayou Grande  
*Enterococcus* CFU/100 ml  
Geomean




Figure 8. Distribution of the geomean of *Enterococcus* counts from Bayou Grande Stations.




Figure 9. Distribution of the maximum of *Enterococcus* counts from Bayou Grande Stations.

Chronic impact areas would be displayed on a plot of minimum *Enterococcus* numbers recorded. For Bayou Grande, none of the minimum recorded values were above the regulatory threshold of a 30 day geomean greater than 35 CFU/100 ml or the 104 CFU/100 ml single sample maximum. The maximum *Enterococcus* numbers (Figure 9) would reflect rain effects on groundwater flow and episodic loading perhaps more relatable to waterfowl or storm water runoff, especially where high maximum values do not correspond to high Geomeans or minimum values. Included in this category would be stations Kalash Dr. storm drain and Payne Rd. storm drain. Highest values of Phosphate and BOD were associated spatially with the high *Enterococcus* geomeans. While variance in the overall dataset prevents these patterns from emerging with traditional statistics, they are apparent visually in the GIS plots.

Analysis of the observed fecal contamination dependence on rainfall provides some additional insights. Analysis of the entire data set (Figure 10), while not providing any statistically significant models, shows high levels of fecal contamination occurring in times of no rainfall. It also indicates that the highest levels of fecal bacteria and the highest variance in system-wide fecal loadings are found following moderate rainfall events (<1”), while a large rainstorm results lower fecal bacteria counts and lower system-wide variance, presumably due to dilution and mixing with storm events.




Figure 10. Bayou Grande *Enterococcus* as a function of rainfall within 48 hours of sampling.

Given the spatially explicit loading patterns found with the GIS analysis and the variance response seen in Figure 10, the dependence of fecal contamination as a function of rainfall was tested for each station. Data associated with rainfall greater than 1.6 “ 48 hours prior to sampling was excluded, as this level of disturbance appears to homogenize the spatial specificity of loadings throughout the system (Figure 10). Two parameters are important outcomes of this analysis. The y-intercept estimate of linear regressions of Log *Enterococcus* as a function of rainfall provides a geomean of *Enterococcus* values for all sampling times with zero rain within the previous 48 hours. This value is indicative of groundwater loading or diffuse, non-storm water sources such as waterfowl. The slope estimates of these functions indicate rainfall influence on *Enterococcus* numbers. Positive slope values indicate increased fecal loading with rainfall as storm water runoff or rain infiltration and enhanced ground water flow. Negative slopes indicate dilution of fecal contamination by rainfall. Because of the high variability in fecal numbers, and the use of these data as relative indicators, a p value of 0.06 was chosen as a threshold of significance for this analysis. Even so, non-significant relationships are still valid as visualization tools in defining trends at the various stations, as outlined below. Some of the variability in counts may be ascribed to tidal mixing and displacement of contamination from its source.

The regression parameters are listed in Table 6. All of the y-intercept estimates are significant, while only two stations' data yielded significant (different from zero) rain effect slope estimates. One of these, station 4, Southside Tributary, had a decline in contamination with increasing rain (negative slope). The other, Station 16 Jamaica Avenue, had contamination significantly increased by moderate rainfall. The lack of other significant effects is due to some combination of high variance, the preponderance of data at zero rainfall, and a real result that rain had little or no impact on contamination levels at these stations.

Despite the low  $R^2$  values, the resulting graphs are valuable as visualization tools, and are included in this report (Figure 11). A high y-intercept and flat (horizontal) or negative slope would be indicative of ground water loading and either a lack of storm water contribution, a storm water contribution that is of the same magnitude (flat) or less (negative by dilution) than the groundwater. A low y-intercept and steep slope would indicate little if any ground water contribution and loading mainly from storm water sources or storm water redistribution of contamination.

The slope values are presented spatially in Figure 12. Geomeans of *Enterococcus* counts for zero rainfall sampling events (y-intercept estimates) are plotted spatially in Figure 13. This GIS analysis clearly shows the impact of contaminated groundwater in the heavily developed area surrounding the creek/ditch stations Fairfield/Rentz Creek, Bartow Avenue, and Paulding Avenue Creek, and the lack of such loading in other areas of the bayou. This analysis provides a better indication of chronically impacted sites than the minimum recorded values from all samples that would be impacted more by mixing and dilution.

Table 6. Regression analysis of Bayou Grande station data as a function of rainfall < 2 inches 48 hours prior to sampling. . Significant parameter estimates ( $p < 0.06$ ) are indicated in bold

| Station Code | Station name                | R <sup>2</sup> | Slope        | p-value      | Intercept    | p-value         | geomean @zero rain |
|--------------|-----------------------------|----------------|--------------|--------------|--------------|-----------------|--------------------|
| 1            | Sherman Grove               | 0.072          | 0.592        | 0.239        | <b>1.485</b> | <b>2.25E-07</b> | <b>30.559</b>      |
| 2            | Kingsport Ave.              | 0.073          | -            | 0.264        | <b>1.483</b> | <b>1.99E-08</b> | <b>30.426</b>      |
| 3            | Acapulco Camino             | 0.110          | 0.695        | 0.142        | <b>1.646</b> | <b>1.60E-08</b> | <b>44.239</b>      |
| 4            | Southside tributary         | 0.139          | -            | 0.096        | <b>1.322</b> | <b>4.36E-08</b> | <b>20.971</b>      |
| 5            | Athens Ave                  | 0.000          | 0.045        | 0.926        | <b>1.037</b> | <b>2.75E-05</b> | <b>10.886</b>      |
| 6            | Inlet at Athens             | 0.022          | 0.315        | 0.523        | <b>1.302</b> | <b>1.36E-06</b> | <b>20.039</b>      |
| 7            | Bremen Ave. tributary       | 0.000          | -            | 0.982        | <b>0.922</b> | <b>5.08E-05</b> | <b>8.358</b>       |
| 8            | Bartow Ave. tributary       | 0.031          | 0.410        | 0.448        | <b>1.232</b> | <b>9.24E-06</b> | <b>17.073</b>      |
| 9            | Cousineau Rd. storm drain   | 0.011          | 0.167        | 0.649        | <b>0.708</b> | <b>7.03E-05</b> | <b>5.106</b>       |
| 10           | Southside site 2            | 0.007          | 0.141        | 0.726        | <b>0.824</b> | <b>3.87E-05</b> | <b>6.672</b>       |
| 11           | Golf Course drain 1         | 0.020          | 0.337        | 0.549        | <b>1.208</b> | <b>2.69E-05</b> | <b>16.127</b>      |
| 12           | Greve Rd. tributary         | 0.074          | 0.511        | 0.246        | <b>1.205</b> | <b>6.99E-07</b> | <b>16.036</b>      |
| 13           | Kalash Dr. storm drain      | 0.001          | 0.067        | 0.910        | <b>0.828</b> | <b>1.55E-03</b> | <b>6.731</b>       |
| 14           | Navy Point bridge (PC15)    | 0.001          | -            | 0.918        | <b>0.991</b> | <b>6.66E-06</b> | <b>9.798</b>       |
| 15           | Oak Ave.                    | 0.099          | 0.656        | 0.165        | <b>1.766</b> | <b>5.33E-09</b> | <b>58.349</b>      |
| 16           | <b>Jamaica Avenue</b>       | <b>0.276</b>   | <b>1.059</b> | <b>0.014</b> | <b>1.362</b> | <b>3.34E-08</b> | <b>23.027</b>      |
| 17           | Syrcler sandbar             | 0.007          | -            | 0.719        | <b>1.234</b> | <b>2.93E-06</b> | <b>17.147</b>      |
| 18           | Syrcler Dr.                 | 0.101          | 0.673        | 0.159        | <b>1.689</b> | <b>1.32E-08</b> | <b>48.858</b>      |
| 19           | Loop Road                   | 0.026          | 0.311        | 0.483        | <b>1.628</b> | <b>9.96E-09</b> | <b>42.501</b>      |
| 20           | Barrios Circle              | 0.003          | 0.087        | 0.821        | <b>1.602</b> | <b>1.53E-09</b> | <b>39.990</b>      |
| 21           | End of bayou                | 0.082          | 0.505        | 0.209        | <b>1.535</b> | <b>4.07E-09</b> | <b>34.310</b>      |
| 22           | Fairfield/Rentz creek       | 0.000          | -            | 0.942        | <b>2.066</b> | <b>3.14E-10</b> | <b>116.416</b>     |
| 23           | Bartow Ave. creek           | 0.013          | 0.199        | 0.679        | <b>2.034</b> | <b>1.35E-07</b> | <b>108.047</b>     |
| 24           | Paulding Ave. creek         | 0.000          | 0.032        | 0.929        | <b>2.404</b> | <b>4.33E-13</b> | <b>253.451</b>     |
| 25           | Baublits Rd. SE storm drain | 0.104          | 0.711        | 0.166        | <b>0.429</b> | <b>3.43E-02</b> | <b>2.687</b>       |
| 26           | Baublits Rd. storm drain    | 0.026          | -            | 0.495        | <b>0.929</b> | <b>2.75E-04</b> | <b>8.493</b>       |
| 27           | Labree Rd. storm drain      | 0.011          | 0.255        | 0.667        | <b>0.870</b> | <b>1.03E-03</b> | <b>7.410</b>       |
| 28           | Midbayou                    | 0.002          | -            | 0.853        | <b>0.622</b> | <b>1.24E-03</b> | <b>4.191</b>       |
| 29           | Navy Point park storm drain | 0.029          | 0.326        | 0.469        | <b>0.465</b> | <b>1.27E-02</b> | <b>2.914</b>       |
| 30           | Palmettos                   | 0.014          | -            | 0.619        | <b>0.718</b> | <b>1.88E-04</b> | <b>5.229</b>       |
| 31           | Payne Rd. storm drain       | 0.056          | 0.596        | 0.316        | <b>0.628</b> | <b>1.05E-02</b> | <b>4.249</b>       |




Figure 11. Station-specific analysis of rainfall dependence in Bayou Grande.




Figure 11. Station-specific analysis of rainfall dependence in Bayou Grande, continued.




Figure 11. Station-specific analysis of rainfall dependence in Bayou Grande, continued.




Figure 11. Station-specific analysis of rainfall dependence in Bayou Grande, continued.




Figure 11. Station-specific analysis of rainfall dependence in Bayou Grande, continued.




Figure 11. Station-specific analysis of rainfall dependence in Bayou Grande, concluded.

# Bayou Grande

## *Enterococcus* Rain Dependence




Figure 12. Slope values from station-specific regression analysis for Bayou Grande




Figure 13. Geomeans of *Enterococcus* at zero rainfall for Bayou Grande estimated as y-intercepts from Regression analysis..

## Bayou Chico

A total of 893 samples were taken over a time period from 14 November 2001 to 30 December 2003. The summary data are presented in Tables 7 & 8.

None of the parameters measured behaved absolutely conservatively, i.e., were diluted from a freshwater source along the salinity gradient in the bayou, although trends are apparent in the data. Figure 14 shows these plots for Nitrate + Nitrite and Phosphate. The nitrogen species appear to be loaded to the system from freshwaters, with a decline towards the bayou mouth and higher salinity. Phosphate, however, shows the opposite trend, suggesting the system may be phosphate limited, and phosphate is drawn into the system from the open bay. Geographic patterns of these data are shown in Figure 15 (Nitrate+Nitrite), Figure 16 (Phosphate), and Figure 17 (biological oxygen demand; BOD).

A conservative mixing analysis of *Enterococcus* counts indicates a stronger freshwater origin for fecal contamination in this system (Figure 18), unlike Bayou Grande, where no relationship of fecal concentrations to salinity was observed.




Figure 14. Conservative mixing diagrams for nitrogen and phosphorous in Bayou Chico.

Table 7. Physico-chemical water quality measures from Bayou Chico sampling.

| site_ID | Site_Description             | Temp  | std  | pH   | std  | Salinity<br>ppt | std  | DO<br>mg/L | std  | BOD<br>mg/L | CV    | NO <sub>3/2</sub><br>µg/L | CV    | TP<br>mg/L | CV    |
|---------|------------------------------|-------|------|------|------|-----------------|------|------------|------|-------------|-------|---------------------------|-------|------------|-------|
| 1       | PYC Boat Ramp                | 21.77 | 6.28 | 7.67 | 0.35 | 17.06           | 7.12 | 7.24       | 1.27 |             |       |                           |       |            |       |
| 2       | Bahia Mar Fuel Dock          | 21.74 | 6.37 | 7.67 | 0.34 | 16.58           | 7.16 | 7.20       | 1.12 |             |       |                           |       |            |       |
| 3       | Pace Storm Drain             | 21.89 | 6.45 | 7.67 | 0.34 | 16.36           | 7.02 | 7.31       | 1.13 |             |       |                           |       |            |       |
| 4       | Runyan's Seawall             | 21.86 | 6.58 | 7.66 | 0.34 | 16.57           | 7.17 | 7.11       | 1.07 |             |       |                           |       |            |       |
| 5       | Scrapyard Fragmites          | 22.12 | 6.43 | 7.61 | 0.42 | 16.12           | 7.21 | 7.29       | 1.05 |             |       |                           |       |            |       |
| 6       | Midbayou (Scrapyard/Island)  | 22.23 | 6.37 | 7.65 | 0.44 | 15.90           | 7.19 | 7.58       | 0.97 |             |       |                           |       |            |       |
| 7       | Pensacola Shipyard A-10      | 22.62 | 6.31 | 7.60 | 0.44 | 16.28           | 7.14 | 7.08       | 1.19 |             |       |                           |       |            |       |
| 8       | Pensacola Shipyard end       | 22.80 | 6.40 | 7.57 | 0.44 | 16.51           | 7.05 | 6.81       | 1.33 |             |       |                           |       |            |       |
| 9       | Tressle Apartments           | 22.56 | 6.37 | 7.47 | 0.41 | 12.82           | 5.97 | 7.96       | 2.03 | 5.14        |       | 111.22                    |       | 0.027      |       |
| 10      | Vince Whibbs GMC Storm Drain | 22.66 | 6.21 | 7.43 | 0.42 | 13.12           | 5.97 | 7.70       | 1.79 |             |       |                           |       |            |       |
| 11      | Navy Boulevard Bridge        | 22.92 | 5.88 | 7.28 | 0.38 | 11.86           | 6.41 | 7.67       | 2.18 | 1.63        | 1.139 | 301.05                    | 0.218 | 0.051      | 0.287 |
| 12      | Church Fragmites             | 23.18 | 5.94 | 7.24 | 0.41 | 12.11           | 6.42 | 7.86       | 2.18 |             |       |                           |       |            |       |
| 13      | Sawgrass at Tin Boat House   | 23.50 | 5.74 | 7.24 | 0.44 | 12.03           | 6.38 | 7.44       | 1.76 | 1.41        | 1.662 | 421.17                    | 0.162 | 0.028      | 0.269 |
| 14      | NE Branch Mouth              | 23.63 | 5.46 | 7.17 | 0.41 | 12.50           | 6.66 | 7.34       | 2.15 |             |       |                           |       |            |       |
| 15      | NE Branch Midway             | 23.41 | 5.82 | 7.00 | 0.37 | 11.25           | 7.35 | 6.27       | 2.09 | 2.39        | 1.024 | 484.15                    | 0.154 | 0.050      | 0.252 |
| 16      | NE Branch East End           | 23.38 | 5.70 | 6.76 | 0.32 | 9.89            | 7.07 | 4.96       | 1.71 |             |       |                           |       |            |       |
| 17      | NW Branch Gazebo             | 23.68 | 5.70 | 7.09 | 0.45 | 10.91           | 5.88 | 7.37       | 2.20 | 2.33        | 0.540 | 614.26                    | 0.154 | 0.040      | 0.192 |
| 18      | NW end                       | 23.54 | 5.42 | 6.99 | 0.41 | 9.48            | 6.23 | 6.36       | 1.66 | 2.33        | 0.615 | 315.29                    | 0.276 | 0.054      | 0.280 |
| 19      | Rip Rap                      | 23.61 | 6.03 | 7.27 | 0.47 | 12.06           | 6.11 | 8.03       | 1.85 | 2.37        | 0.568 | 362.71                    | 0.137 | 0.032      | 0.150 |
| 20      | Juncus at Apartments         | 23.21 | 6.17 | 7.38 | 0.45 | 12.29           | 6.09 | 8.10       | 2.08 | 1.77        | 0.957 | 244.46                    | 0.232 | 0.047      | 0.257 |
| 21      | Channel Marker 17            | 23.13 | 6.58 | 7.68 | 0.49 | 16.25           | 6.93 | 7.96       | 1.05 | 1.23        | 3.889 | 73.91                     | 0.125 | 0.067      | 0.302 |
| 22      | Rope Fence                   | 23.08 | 6.16 | 7.52 | 0.56 | 14.60           | 7.76 | 7.50       | 1.65 |             |       |                           |       |            |       |
| 23      | Lakewood Park                | 22.93 | 6.41 | 7.55 | 0.58 | 14.65           | 7.43 | 7.74       | 1.31 |             |       |                           |       |            |       |
| 24      | West Branch Cattails         | 23.42 | 6.15 | 7.42 | 0.55 | 14.37           | 8.07 | 7.20       | 1.48 |             |       |                           |       |            |       |
| 25      | West Branch Marsh Point      | 23.28 | 5.80 | 7.24 | 0.61 | 10.94           | 7.24 | 6.54       | 1.83 |             |       | 426.40                    | 0.028 | 0.032      | 0.281 |
| 26      | West end Last Dock           | 25.19 | 4.35 | 7.13 | 0.65 | 10.52           | 6.33 | 6.07       | 1.81 | 2.32        | 0.373 | 72.04                     | 0.409 | 0.055      | 0.355 |
| 27      | Swamplillies at Green Roof   | 23.35 | 6.52 | 7.46 | 0.68 | 14.05           | 7.76 | 7.67       | 1.54 |             |       |                           |       |            |       |
| 28      | Tire Pole                    | 23.45 | 6.65 | 7.55 | 0.51 | 15.40           | 7.41 | 7.61       | 1.18 |             |       |                           |       |            |       |
| 29      | Bell Marine Fragmites        | 22.52 | 6.83 | 7.62 | 0.46 | 15.42           | 7.24 | 7.83       | 1.09 |             |       |                           |       |            |       |
| 30      | Pelican Pole                 | 22.56 | 6.79 | 7.67 | 0.42 | 16.06           | 7.17 | 7.60       | 1.02 |             |       |                           |       |            |       |
| 31      | Mahogany Landing             | 22.24 | 6.96 | 7.66 | 0.38 | 16.00           | 7.00 | 7.47       | 1.13 | 1.29        | 2.009 | 53.02                     | 0.418 | 0.051      | 0.295 |
| 32      | Marker 10/ Pilings           | 22.26 | 6.59 | 7.78 | 0.34 | 17.40           | 7.29 | 7.54       | 1.41 |             |       |                           |       |            |       |
| 33      | Ditch                        | 22.13 | 4.07 | 7.33 | 0.55 | 7.59            | 9.81 | 6.86       | 1.57 | 2.28        | 0.500 | 1760.21                   | 0.061 | 0.030      | 0.000 |
| 34      | S-Street                     | 24.21 | 4.06 | 6.64 | 0.31 | 0.00            | 0.00 | 5.22       | 2.30 | 3.66        | 0.462 | 549.71                    | 0.100 | 0.037      | 0.244 |
| 35      | Corry Field Road North       | 22.17 | 4.03 | 6.93 | 0.40 | 0.05            | 0.05 | 5.49       | 1.67 | 1.81        |       | 1314.40                   |       | 0.030      |       |
| 36      | Corry Field Road South       | 21.97 | 5.07 | 6.42 | 0.45 | 2.32            | 4.78 | 4.84       | 1.56 | 1.88        | 0.113 | 202.62                    | 0.095 | 0.026      | 0.102 |
| 37      | Brigadier                    | 21.66 | 4.97 | 5.71 | 0.56 | 0.00            | 0.00 | 4.50       | 1.57 | 1.20        | 5.326 | 39.05                     | 0.272 | 0.032      | 0.426 |

Table 7, continued. Physico-chemical water quality measures from Bayou Chico sampling.

| site_ID | Site_Description    | Temp  | std  | pH   | std  | Salinity<br>ppt | std  | DO<br>mg/L | std  | BOD<br>mg/L | CV    | NO <sub>3/2</sub><br>µg/L | CV    | TP<br>mg/L | CV    |
|---------|---------------------|-------|------|------|------|-----------------|------|------------|------|-------------|-------|---------------------------|-------|------------|-------|
| 38      | Fairfield           | 20.91 | 5.79 | 5.11 | 1.15 | 0.00            | 0.00 | 2.88       | 2.03 | 1.79        | 0.145 | 6.77                      | 0.409 | 0.024      | 0.213 |
| 39      | Q-Street            | 24.62 | 2.47 | 6.71 | 0.07 | 0.00            | 0.00 | 3.63       | 1.06 | 0.85        | 6.610 | 1331.82                   | 0.012 | 0.030      | 0.000 |
| 40      | New Warrington      | 23.15 | 3.30 | 6.51 | 0.56 | 0.05            | 0.10 | 5.01       | 1.13 | 3.35        | 0.615 | 224.42                    | 0.044 | 0.036      | 0.128 |
| 41      | Twin Oaks Apartment | 21.66 | 5.05 | 6.92 | 0.28 | 0.00            | 0.00 | 6.41       | 2.83 | 2.17        | 0.405 | 1441.27                   | 0.016 | 0.041      | 0.185 |
| 42      | Twin Oaks/Prieto    | 20.46 | 4.71 | 6.99 | 0.11 | 0.00            | 0.00 | 8.15       | 2.02 | 1.81        |       | 2252.10                   |       | 0.030      |       |

Table 8. Summary fecal indicator data from Bayou Chico sampling.

| site_ID | Site_Description             | Geomean Entero | CV    | entero min | entero max |
|---------|------------------------------|----------------|-------|------------|------------|
| 1       | PYC Boat Ramp                | 23.58          | 0.427 | 2          | 510        |
| 2       | Bahia Mar Fuel Dock          | 26.30          | 0.426 | 4          | 640        |
| 3       | Pace Storm Drain             | 34.68          | 0.411 | 2          | 400        |
| 4       | Runyan's Seawall             | 12.59          | 0.544 | 2          | 270        |
| 5       | Scrapyard Fragmites          | 21.79          | 0.444 | 2          | 450        |
| 6       | Midbayou (Scrapyard/Island)  | 14.56          | 0.478 | 2          | 310        |
| 7       | Pensacola Shipyard A-10      | 14.54          | 0.561 | 2          | 950        |
| 8       | Pensacola Shipyard end       | 16.27          | 0.402 | 2          | 132        |
| 9       | Tressle Apartments           | 64.90          | 0.302 | 14         | 940        |
| 10      | Vince Whibbs GMC Storm Drain | 66.18          | 0.329 | 6          | 1440       |
| 11      | Navy Boulevard Bridge        | 93.05          | 0.379 | 10         | 4080       |
| 12      | Church Fragmites             | 88.64          | 0.340 | 20         | 4260       |
| 13      | Sawgrass at Tin Boat House   | 102.46         | 0.363 | 8          | 2600       |
| 14      | NE Branch Mouth              | 96.29          | 0.333 | 6          | 5220       |
| 15      | NE Branch Midway             | 144.02         | 0.322 | 10         | 6790       |
| 16      | NE Branch East End           | 146.76         | 0.288 | 22         | 4370       |
| 17      | NW Branch Gazebo             | 116.81         | 0.289 | 8          | 6380       |
| 18      | NW end                       | 146.47         | 0.292 | 16         | 5460       |
| 19      | Rip Rap                      | 45.69          | 0.442 | 2          | 2310       |
| 20      | Juncus at Apartments         | 24.50          | 0.505 | 2          | 1020       |
| 21      | Channel Marker 17            | 11.28          | 0.611 | 2          | 1080       |
| 22      | Rope Fence                   | 18.25          | 0.492 | 2          | 320        |
| 23      | Lakewood Park                | 14.65          | 0.439 | 2          | 202        |
| 24      | West Branch Cattails         | 20.16          | 0.396 | 2          | 126        |
| 25      | West Branch Marsh Point      | 58.19          | 0.366 | 2          | 1130       |
| 26      | West end Last Dock           | 59.32          | 0.325 | 4          | 790        |
| 27      | Swamplillies at Green Roof   | 27.49          | 0.419 | 2          | 600        |
| 28      | Tire Pole                    | 28.44          | 0.381 | 2          | 329        |
| 29      | Bell Marine Fragmites        | 24.35          | 0.436 | 2          | 280        |
| 30      | Pelican Pole                 | 22.38          | 0.473 | 2          | 271        |
| 31      | Mahogany Landing             | 20.62          | 0.453 | 2          | 220        |
| 32      | Marker 10/ Pilings           | 11.59          | 0.625 | 2          | 580        |
| 33      | Ditch                        | 201.53         | 0.305 | 2          | 3990       |
| 34      | S-Street                     | 429.14         | 0.136 | 74         | 1260       |
| 35      | Corry Field Road North       | 407.54         | 0.203 | 0          | 2950       |
| 36      | Corry Field Road South       | 179.92         | 0.211 | 8          | 2400       |
| 37      | Brigadier                    | 159.39         | 0.225 | 10         | 1380       |
| 38      | Fairfield                    | 45.72          | 0.310 | 4          | 600        |
| 39      | Q-Street                     | 329.76         | 0.162 | 80         | 1010       |
| 40      | New Warrington               | 156.73         | 0.209 | 58         | 450        |
| 41      | Twin Oaks Apartment          | 445.82         | 0.094 | 240        | 960        |
| 42      | Twin Oaks/Prieto             | 376.29         | 0.110 | 132        | 600        |

## Bayou Chico Nitrate + Nitrite (ug/L)




Figure 15. Mean Nitrate+Nitrite values for Bayou Chico station data.

# Bayou Chico Total Phosphorous (mg/L)




Figure 16. Mean Phosphate values for Bayou Chico station data.

## Bayou Chico Biological Oxygen Demand (mg/L)




Figure 17. Mean BOD values for Bayou Chico station data.




Figure 18. Conservative mixing analysis of *Enterococcus* in Bayou Chico.

Geospatial analysis of these data by station add resolution to the pattern of higher contamination levels at lower salinities. Figure 19 displays the distribution of contamination as the geomean of each station's data. With the exception of the drainage stream at the mouth of the bayou, the only stations with geomeans exceeding the 30 day geomean regulatory limit of 35 CFU/100 ml sample were located in the upper freshwater reaches of the bayou. This distribution is reinforced by the plot of minimum *Enterococcus* counts recorded at each station (Figure 20), which would be indicative of chronically impaired stations. Maximum recorded counts by station (Figure 21) should reflect storm water effects as well as chronic effects, and a few stations in the main part of the bayou reflect these episodic loadings. However the predominant pattern for the maximum recorded counts follows the patterns for geomeans and minimum counts.

## Bayou Chico *Enterococcus* CFU/100 ml Geomean




Figure 19. Geomeans of Bayou Chico station data for *Enterococcus* counts.

Bayou Chico  
*Enterococcus* CFU/100 ml  
 Minimum Count




Figure 20. Bayou Chico station data for *Enterococcus* minimum counts.

Bayou Chico  
*Enterococcus* CFU/100 ml  
Maximum Count




Figure 21. Bayou Chico station data for *Enterococcus* maximum counts.

Using correlation analysis, none of the parameters measured were predictive of fecal bacteria concentrations within the system. The highest correlation found with rainfall data was for *Enterococcus* and rain total on the day of sampling (0.23). Correlations with the previous day rainfall and the sum of rainfall for up to 7 days prior to sampling were much lower (data not shown).

Analysis of the entire *Enterococcus* data against the amount of rainfall in the previous 48 hours shows a pattern similar to that found in Bayou Grande, yet more pronounced, with the highest recorded counts occurring during moderate rainfall (0.5-1.0") and lower counts obtained following heavier rainfall (Figure 22). Regression analysis of the *Enterococcus* count dependence on rainfall by station is shown in Figure 23 and summarized in Table 9. Stations 39-42 were excluded from this analysis due to too few data points (4-5). The data for a majority of stations returned significant slope and intercept estimates. Those with insignificant slope estimates tended to have high intercept values. Indeed, these parameters were negatively correlated (-0.732). The analysis indicates strong rain dependence for the *Enterococcus* loading into the lower portion of Bayou Chico, and chronic loading in fresher upper reaches. Slope and intercept estimates are shown plotted in GIS analysis in Figure 24 (slopes) and Figure 25 (intercepts).




Figure 22. Bayou Chico *Enterococcus* as a function of rainfall in the 48 hours prior to sampling.

Table 9. Regression of Bayou Chico station data as a function of rainfall < 2 inches 48 hours prior to sampling. Significant parameter estimates ( $p < 0.06$ ) are indicated in bold.

| Site ID | Site Description                    | R <sup>2</sup> | Slope        | p value      | Intercept    | p value         | Geomean @ zero rain |
|---------|-------------------------------------|----------------|--------------|--------------|--------------|-----------------|---------------------|
| 1       | PYC Boat Ramp                       | 0.135          | 0.410        | 0.122        | <b>1.184</b> | <b>1.85E-07</b> | <b>15.29</b>        |
| 2       | <b>Bahia Mar Fuel Dock</b>          | <b>0.297</b>   | <b>0.628</b> | <b>0.016</b> | <b>1.160</b> | <b>9.07E-08</b> | <b>14.44</b>        |
| 3       | <b>Pace Storm Drain</b>             | <b>0.204</b>   | <b>0.569</b> | <b>0.052</b> | <b>1.308</b> | <b>1.42E-07</b> | <b>20.34</b>        |
| 4       | Runyan's Seawall                    | 0.092          | 0.332        | 0.206        | <b>0.946</b> | <b>3.87E-06</b> | <b>8.83</b>         |
| 5       | <b>Scrapyard Fragmites</b>          | <b>0.205</b>   | <b>0.481</b> | <b>0.052</b> | <b>1.117</b> | <b>1.20E-07</b> | <b>13.08</b>        |
| 6       | <b>Midbayou (Scrapyard/Island)</b>  | <b>0.368</b>   | <b>0.608</b> | <b>0.006</b> | <b>0.920</b> | <b>1.58E-07</b> | <b>8.32</b>         |
| 7       | Pensacola Shipyard A-10             | 0.018          | 0.176        | 0.584        | <b>1.066</b> | <b>1.32E-05</b> | <b>11.65</b>        |
| 8       | Pensacola Shipyard end              | 0.024          | 0.154        | 0.522        | <b>1.141</b> | <b>1.28E-07</b> | <b>13.83</b>        |
| 9       | <b>Tressle Apartments</b>           | <b>0.323</b>   | <b>0.644</b> | <b>0.004</b> | <b>1.569</b> | <b>1.31E-12</b> | <b>37.04</b>        |
| 10      | <b>Vince Whibbs GMC Storm Drain</b> | <b>0.247</b>   | <b>0.633</b> | <b>0.014</b> | <b>1.580</b> | <b>3.15E-11</b> | <b>38.00</b>        |
| 11      | <b>Navy Boulevard Bridge</b>        | <b>0.339</b>   | <b>0.945</b> | <b>0.003</b> | <b>1.641</b> | <b>4.40E-10</b> | <b>43.71</b>        |
| 12      | <b>Church Fragmites</b>             | <b>0.174</b>   | <b>0.587</b> | <b>0.043</b> | <b>1.724</b> | <b>9.72E-11</b> | <b>52.98</b>        |
| 13      | <b>Sawgrass at Tin Boat House</b>   | <b>0.307</b>   | <b>0.884</b> | <b>0.005</b> | <b>1.712</b> | <b>2.21E-10</b> | <b>51.58</b>        |
| 14      | <b>NE Branch Mouth</b>              | <b>0.312</b>   | <b>0.816</b> | <b>0.005</b> | <b>1.708</b> | <b>4.12E-11</b> | <b>51.07</b>        |
| 15      | <b>NE Branch Midway</b>             | <b>0.341</b>   | <b>0.895</b> | <b>0.003</b> | <b>1.855</b> | <b>1.38E-11</b> | <b>71.66</b>        |
| 16      | <b>NE Branch East End</b>           | <b>0.169</b>   | <b>0.565</b> | <b>0.046</b> | <b>1.961</b> | <b>5.47E-12</b> | <b>91.50</b>        |
| 17      | <b>NW Branch Gazebo</b>             | <b>0.167</b>   | <b>0.541</b> | <b>0.047</b> | <b>1.897</b> | <b>5.03E-12</b> | <b>78.92</b>        |
| 18      | <b>NW end</b>                       | <b>0.241</b>   | <b>0.698</b> | <b>0.015</b> | <b>1.965</b> | <b>4.12E-12</b> | <b>92.28</b>        |
| 19      | <b>Rip Rap</b>                      | <b>0.322</b>   | <b>0.894</b> | <b>0.004</b> | <b>1.316</b> | <b>1.64E-08</b> | <b>20.71</b>        |
| 20      | <b>Juncus at Apartments</b>         | <b>0.263</b>   | <b>0.747</b> | <b>0.010</b> | <b>1.124</b> | <b>1.26E-07</b> | <b>13.32</b>        |
| 21      | <b>Channel Marker 17</b>            | <b>0.231</b>   | <b>0.625</b> | <b>0.037</b> | <b>0.832</b> | <b>5.09E-05</b> | <b>6.79</b>         |
| 22      | Rope Fence                          | 0.064          | 0.318        | 0.297        | <b>1.133</b> | <b>2.88E-06</b> | <b>13.57</b>        |
| 23      | Lakewood Park                       | 0.163          | 0.395        | 0.087        | <b>1.011</b> | <b>2.20E-07</b> | <b>10.25</b>        |
| 24      | West Branch Cattails                | 0.017          | 0.145        | 0.593        | <b>1.258</b> | <b>1.80E-07</b> | <b>18.13</b>        |
| 25      | West Branch Marsh Point             | 0.087          | 0.419        | 0.183        | <b>1.625</b> | <b>1.07E-08</b> | <b>42.18</b>        |
| 26      | West end Last Dock                  | 0.086          | 0.316        | 0.289        | <b>1.654</b> | <b>6.67E-07</b> | <b>45.05</b>        |
| 27      | Swampillies at Green Roof           | 0.116          | 0.436        | 0.154        | <b>1.288</b> | <b>4.49E-07</b> | <b>19.39</b>        |
| 28      | <b>Tire Pole</b>                    | <b>0.259</b>   | <b>0.576</b> | <b>0.026</b> | <b>1.243</b> | <b>3.88E-08</b> | <b>17.48</b>        |
| 29      | <b>Bell Marine Fragmites</b>        | <b>0.230</b>   | <b>0.597</b> | <b>0.038</b> | <b>1.170</b> | <b>4.39E-07</b> | <b>14.80</b>        |
| 30      | <b>Pelican Pole</b>                 | <b>0.255</b>   | <b>0.652</b> | <b>0.027</b> | <b>1.108</b> | <b>1.21E-06</b> | <b>12.82</b>        |
| 31      | Mahogany Landing                    | 0.161          | 0.498        | 0.089        | <b>1.144</b> | <b>1.02E-06</b> | <b>13.93</b>        |
| 32      | Marker 10/ Pilings                  | 0.039          | 0.245        | 0.417        | <b>0.930</b> | <b>2.99E-05</b> | <b>8.50</b>         |
| 33      | Ditch                               | 0.000          | 0.002        | 0.994        | <b>2.253</b> | <b>1.45E-11</b> | <b>178.92</b>       |
| 34      | S-Street                            | 0.001          | 0.019        | 0.912        | <b>2.611</b> | <b>6.65E-19</b> | <b>408.78</b>       |
| 35      | <b>Corry Field Road North</b>       | <b>0.368</b>   | <b>0.612</b> | <b>0.006</b> | <b>2.373</b> | <b>3.81E-13</b> | <b>236.01</b>       |
| 36      | Corry Field Road South              | 0.143          | 0.396        | 0.069        | <b>2.152</b> | <b>4.69E-15</b> | <b>141.94</b>       |
| 37      | Brigadier                           | 0.042          | 0.225        | 0.385        | <b>2.167</b> | <b>5.06E-12</b> | <b>146.95</b>       |
| 38      | Fairfield                           | 0.106          | 0.380        | 0.121        | <b>1.536</b> | <b>5.21E-11</b> | <b>34.35</b>        |









Figure 23. *Enterococcus* dependence on rainfall by station for Bayou Chico.









Figure 23, continued. *Enterococcus* dependence on rainfall by station for Bayou Chico.









Figure 23, continued. *Enterococcus* dependence on rainfall by station for Bayou Chico.









Figure 23, continued. *Enterococcus* dependence on rainfall by station for Bayou Chico.









Figure 23, continued. *Enterococcus* dependence on rainfall by station for Bayou Chico.





Figure 23, concluded. *Enterococcus* dependence on rainfall by station for Bayou Chico.

# Bayou Chico

## Enterococcus Rain Dependence




Figure 24. Distribution of slope values for Bayou Chico station rain dependence.

Bayou Chico  
*Enterococcus* CFU/100 ml  
 Geomean at Zero Rainfall




Figure 25. Distribution of y-intercept values (geomeans at zero rainfall) for Bayou Chico Stations..

## Bayou Texar

A total of 893 samples were taken over a time period from 14 November 2001 to 30 December 2003. The summary data are presented in Tables 10 and 11.

A conservative mixing analysis for Bayou Texar nutrient data is shown in Figure 26. Nitrate+nitrite and phosphate show similar patterns as Bayou Chico, with the nitrogen sources having more of a freshwater origin and the phosphate tending to have an open bay source. However, the trend was not as clear nor were nitrogen concentrations as high as those recorded for Bayou Chico.




Figure 26. Conservative mixing diagrams for Nitrate + Nitrite and Phosphate in Bayou Texar.

GIS maps for the distribution of Nitrogen, Phosphate and Biological Oxygen Demand (BOD) are shown in Figures 27, 28, & 29.

Table 10. Physico-chemical water quality measures from Bayou Texar sampling.

| station code | station name              | Temp   | std   | Ph    | std   | salinity | std   | DO mg/L | std   | BOD mg/L | CV    | NO <sub>3/2</sub> µg/L | CV    | TP mg/L | CV    |
|--------------|---------------------------|--------|-------|-------|-------|----------|-------|---------|-------|----------|-------|------------------------|-------|---------|-------|
| 1            | Cervantes Bridge          | 21.543 | 6.618 | 7.645 | 0.429 | 16.675   | 6.266 | 7.491   | 2.033 | 1.360    |       | 200                    |       | 0.050   |       |
| 2            | Brainerd St. pond         | 21.757 | 6.422 | 7.565 | 0.552 | 15.577   | 6.688 | 6.482   | 2.230 | 1.360    |       | 100                    |       | 0.100   |       |
| 3            | Bayview Park pvc SD       | 21.484 | 6.615 | 7.709 | 0.283 | 17.444   | 5.746 | 7.587   | 1.762 | 1.510    |       | 200                    |       | 0.080   |       |
| 4            | Tree Roots                | 21.331 | 6.724 | 7.710 | 0.281 | 18.428   | 5.110 | 7.742   | 1.631 |          |       |                        |       |         |       |
| 5            | Boathouse (point)         | 22.123 | 6.778 | 7.703 | 0.350 | 16.253   | 5.715 | 7.339   | 2.187 |          |       |                        |       |         |       |
| 6            | Rocks/Gazebo              | 21.273 | 6.622 | 7.672 | 0.364 | 17.788   | 5.537 | 7.942   | 1.873 |          |       |                        |       |         |       |
| 7            | Oriental Garden           | 21.422 | 6.596 | 7.637 | 0.357 | 17.292   | 5.761 | 7.886   | 1.759 |          |       |                        |       |         |       |
| 8            | South Whaley fragmites    | 21.284 | 6.488 | 7.629 | 0.360 | 17.488   | 5.769 | 7.843   | 1.512 | 1.380    |       | 200                    |       | 0.080   |       |
| 9            | Whaley Ditch storm drain  | 22.106 | 6.670 | 7.767 | 1.003 | 15.322   | 6.190 | 7.676   | 1.845 | 1.240    |       | 200                    |       | 0.090   |       |
| 10           | Birnam Woods green SD     | 21.642 | 6.438 | 7.650 | 0.365 | 17.232   | 5.785 | 8.144   | 1.466 |          |       |                        |       |         |       |
| 11           | Blackshear Ave. SD        | 22.597 | 6.587 | 7.450 | 0.297 | 14.195   | 6.185 | 7.356   | 1.847 | 1.297    | 0.098 | 170.71                 | 0.003 | 0.080   | 0.332 |
| 12           | Blanford place FW seep    | 22.748 | 6.691 | 7.203 | 0.300 | 11.733   | 6.610 | 7.258   | 1.811 |          |       |                        |       |         |       |
| 13           | 34th St. storm drain      | 22.308 | 6.302 | 7.100 | 0.312 | 12.196   | 7.325 | 7.512   | 1.534 | 1.207    | 0.157 | 422.26                 | 0.001 | 0.112   | 0.221 |
| 14           | Six Cement poles          | 23.082 | 6.009 | 7.043 | 0.357 | 12.496   | 7.170 | 7.562   | 1.684 |          |       |                        |       |         |       |
| 15           | Carpenter Creek center    | 22.963 | 5.643 | 6.888 | 0.316 | 9.960    | 7.184 | 7.242   | 1.042 | 0.860    |       | 900                    |       | 0.070   |       |
| 16           | Driftwood 4 SD            | 23.117 | 6.425 | 6.896 | 0.290 | 10.798   | 7.033 | 6.514   | 1.918 | 1.467    | 0.186 | 470.59                 | 0.001 | 0.078   | 0.322 |
| 17           | Texar Woods SD            | 23.254 | 6.290 | 7.272 | 0.408 | 13.745   | 6.446 | 7.474   | 1.973 |          |       |                        |       |         |       |
| 18           | Seville Dr. (2) SD        | 22.901 | 6.406 | 7.352 | 0.345 | 14.346   | 6.065 | 7.048   | 1.931 | 1.360    |       | 400                    |       | 0.080   |       |
| 19           | Banquos Court SD          | 22.661 | 6.609 | 7.510 | 0.419 | 14.320   | 6.523 | 7.532   | 1.835 | 1.472    | 0.117 | 113.44                 | 0.005 | 0.089   | 0.491 |
| 20           | Bayou Blvd./Perry SD      | 22.429 | 6.825 | 7.687 | 0.320 | 15.938   | 5.877 | 7.671   | 1.968 |          |       |                        |       |         |       |
| 21           | 12th Ave. bridge          | 21.939 | 4.323 | 6.768 | 0.401 | 1.681    | 3.480 | 6.379   | 1.736 | 1.134    | 0.115 | 1247.40                | 0.000 | 0.031   | 0.310 |
| 22           | 9th Ave.                  | 21.072 | 3.064 | 6.649 | 0.435 | 0.000    | 0.000 | 7.434   | 1.559 | 2.428    | 0.268 | 1.22                   | 0.121 | 0.332   |       |
| 23           | Airport Blvd.             | 20.861 | 5.607 | 6.697 | 0.347 | 0.000    | 0.000 | 7.290   | 2.020 | 1.340    | 0.158 | 260.10                 | 0.002 | 0.088   | 0.096 |
| 24           | Born Court                | 19.650 | 5.210 | 6.561 | 0.699 | 0.004    | 0.021 | 5.608   | 1.875 | 1.742    | 0.035 | 332.95                 | 0.001 | 0.031   | 0.356 |
| 25           | Boiling Brook             | 20.181 | 5.386 | 6.733 | 0.553 | 0.002    | 0.015 | 6.142   | 1.783 | 1.893    | 0.105 | 1088.64                | 0.000 | 0.040   | 0.225 |
| 26           | Sears Warehouse           | 18.789 | 4.574 | 6.092 | 0.917 | 0.000    | 0.000 | 2.495   | 1.444 | 2.209    | 0.078 | 33.34                  | 0.003 | 0.017   | 0.348 |
| 27           | Interstate 10-Hist. Dist. | 21.395 | 6.359 | 6.696 | 0.343 | 0.005    | 0.022 | 5.233   | 2.561 |          |       |                        |       |         |       |
| 28           | Olive Road                | 22.392 | 6.523 | 6.569 | 0.323 | 0.000    | 0.000 | 4.046   | 2.606 | 1.994    | 0.027 | 17.89                  | 0.017 | 0.027   | 0.224 |
| 29           | Walton/Davis              | 20.456 | 6.491 | 6.842 | 0.512 | 0.000    | 0.000 | 4.768   | 2.657 | 0.865    | 0.220 | 72.50                  | 0.008 |         |       |
| 30           | Brookside Place           | 21.213 | 3.808 | 6.405 | 0.200 | 0.000    | 0.000 | 5.815   | 1.416 | 1.277    | 0.086 |                        |       | 0.005   | 0.000 |
| 31           | Creekside Office          | 22.731 | 7.591 | 6.723 | 0.102 | 0.000    | 0.000 | 4.362   | 1.559 | 1.839    | 0.142 | 6.84                   | 0.042 | 0.055   | 0.369 |
| 32           | Springhill                | 21.296 | 4.063 | 6.528 | 0.079 | 0.000    | 0.000 | 7.408   | 1.331 | 0.401    | 0.351 | 1013.77                | 0.000 |         |       |
| 33           | Burgess Road              | 21.511 | 4.525 | 7.168 | 0.399 | 0.000    | 0.000 | 5.954   | 0.601 | 0.327    | 0.548 | 199.43                 | 0.000 | 0.008   | 0.624 |

Table 11. Summary fecal indicator data from Bayou Texar sampling.

| station code | station name                   | Entero geomean | CV    | min | max   | Geomean FC | CV    | min | max   |
|--------------|--------------------------------|----------------|-------|-----|-------|------------|-------|-----|-------|
| 1            | Cervantes Bridge               | 15.221         | 0.450 | 1   | 1140  | 29.008     | 0.224 | 20  | 500   |
| 2            | Brainerd St. pond              | 79.743         | 0.544 | 1   | 16000 | 427.820    | 0.371 | 20  | 16000 |
| 3            | Bayview Park pvc/storm drain   | 9.647          | 0.447 | 1   | 172   | 40.000     |       | 40  | 40    |
| 4            | Tree Roots                     | 4.650          | 0.587 | 1   | 120   |            |       |     |       |
| 5            | Boathouse (point)              | 6.745          | 0.525 | 1   | 103   |            |       |     |       |
| 6            | Rocks/Gazebo                   | 4.963          | 0.597 | 1   | 188   |            |       |     |       |
| 7            | Oriental Garden                | 5.224          | 0.596 | 1   | 240   |            |       |     |       |
| 8            | South Whaley fragmites         | 5.340          | 0.612 | 1   | 380   | 20.000     |       | 20  | 20    |
| 9            | Whaley Ditch storm drain       | 11.396         | 0.495 | 1   | 408   | 40.000     |       | 40  | 40    |
| 10           | Birnam Woods green SD          | 18.988         | 0.402 | 1   | 600   | 51.837     | 0.363 | 20  | 5000  |
| 11           | Blackshear Ave. SD             | 40.575         | 0.349 | 2   | 1770  | 159.933    | 0.345 | 20  | 16000 |
| 12           | Blanford place FW seep         | 55.444         | 0.308 | 8   | 1240  | 167.949    | 0.272 | 20  | 3000  |
| 13           | 34th St. storm drain           | 60.579         | 0.336 | 11  | 1180  | 219.551    | 0.254 | 20  | 3000  |
| 14           | Six Cement poles-tan house     | 57.998         | 0.328 | 6   | 1950  | 161.002    | 0.323 | 20  | 2400  |
| 15           | Carpenter Creek center         | 115.489        | 0.311 | 7   | 1990  | 417.943    | 0.220 | 40  | 16000 |
| 16           | Driftwood 4 SD                 | 105.768        | 0.307 | 5   | 2200  | 343.619    | 0.241 | 20  | 3000  |
| 17           | Texar Woods SD                 | 34.677         | 0.346 | 2   | 600   | 104.168    | 0.319 | 20  | 2200  |
| 18           | Seville Dr. (2) SD             | 68.320         | 0.297 | 7   | 660   | 151.024    | 0.343 | 20  | 9000  |
| 19           | Banquos Court SD               | 55.676         | 0.510 | 1   | 16000 | 398.377    | 0.401 | 20  | 16000 |
| 20           | Bayou Blvd./Perry SD           | 20.915         | 0.501 | 1   | 3010  | 58.954     | 0.391 | 20  | 5000  |
| 21           | 12th Ave. bridge               | 283.488        | 0.236 | 18  | 5000  | 783.117    | 0.259 | 20  | 16000 |
| 22           | 9th Ave.                       | 326.239        | 0.184 | 34  | 11800 | 313.807    | 0.240 | 20  | 5000  |
| 23           | Airport Blvd.                  | 187.684        | 0.222 | 20  | 1100  | 279.237    | 0.310 | 20  | 16000 |
| 24           | Born Court                     | 308.484        | 0.250 | 4   | 2220  | 214.076    | 0.265 | 20  | 5000  |
| 25           | Boiling Brook                  | 299.474        | 0.225 | 18  | 3120  | 147.352    | 0.283 | 20  | 1700  |
| 26           | Sears Warehouse                | 108.716        | 0.381 | 1   | 3350  | 421.949    | 0.452 | 20  | 16000 |
| 27           | Interstate 10-Historical Dist. | 151.069        | 0.269 | 4   | 1700  | 218.682    | 0.422 | 20  | 30000 |
| 28           | Olive Road                     | 88.224         | 0.325 | 1   | 1500  | 171.900    | 0.290 | 40  | 3000  |
| 29           | Walton/Davis                   | 548.073        | 0.208 | 18  | 6800  |            |       |     |       |
| 30           | Brookside Place                | 495.211        | 0.066 | 260 | 940   |            |       |     |       |
| 31           | Creekside Office               | 60.371         | 0.315 | 20  | 680   |            |       |     |       |
| 32           | Springhill                     | 475.892        | 0.143 | 83  | 2360  |            |       |     |       |
| 33           | Burgess Road                   | 575.627        | 0.108 | 200 | 1340  |            |       |     |       |

# Bayou Texar Nitrate + Nitrite (ug/L)




Figure 27. Distribution of mean Nitrate+Nitrite values in samples from Bayou Texar stations.

# Bayou Texar Total Phosphorous (mg/L)




Figure 28. Mean Phosphate values in samples from Bayou Texar stations.

# Bayou Texar Biological Oxygen Demand (mg/L)




Figure 29. Mean BOD values in samples from Bayou Texar stations.

*Enterococcus* counts as a function of salinity indicate a predominantly freshwater source for fecal contamination within the saline portions of Bayou Texar (Figure 30), although considerable variation is apparent.




Figure 30. Conservative mixing diagrams for *Enterococcus* in Bayou Texar, including only stations south from the 12 Avenue Bridge.

Geomeans for each station's data is plotted by GIS in Figure 31. These data clearly show the upper reaches of the bayou and Carpenter's Creek Stations as dominating the loading of fecal material. This pattern is further accentuated by examining the chronically loaded stations with the plot of minimum *Enterococcus* counts (Figure 32). Maximum counts recorded show the influence of storm water runoff in the bayou (Figure 33), with the highest counts occurring in the main part of the bayou as well as up into Carpenters Creek.

Bayou Texar  
*Enterococcus* CFU/100 ml  
Geomean




Figure 31. Geomeans for *Enterococcus* counts at each station in Bayou Texar.

Bayou Texar  
*Enterococcus* CFU/100 ml  
Minimum Count




Figure 32. Minimum *Enterococcus* counts recorded from each station in Bayou Texar.

Bayou Texar  
*Enterococcus* CFU/100 ml  
Maximum Count




Figure 33. Maximum *Enterococcus* counts recorded from each station in Bayou Texar.

Using correlation analysis, none of the parameters measured were predictive of fecal bacteria loadings within the system. The highest correlation found with rainfall data was for fecal coliforms and the rain total for the 48 hours prior to sampling (0.316). By regression analysis (Figure 34), Bayou Texar had the clearest rain influence of all three bayous. The pattern of reduced variance in *Enterococcus* numbers with high rainfall seen in the other bayous was also found in Bayou Texar, but the reduction of variance was due more to the loss of low counts within the system than decreased high counts from dilution (Figure 34). The impact of rain on Bayou Texar was also reflected in the number of stations with data yielding significant slope estimates for *Enterococcus* dependence on rainfall (Table 12; Figure 35). As with Chico, these slope and intercept estimates were negatively correlated (-0.587), suggesting some separation between groundwater and storm water influences by station. The geospatial distribution of the slope estimates indicates fecal loadings associated with rainfall and storm water runoff throughout the system (Figure 36), but the chronically impaired stations showing high *Enterococcus* counts without any rain prior to sampling as occurring within Carpenters Creek and not in the main part of Bayou Texar (Figure 37)




Figure 34. *Enterococcus* counts from Bayou Texar as a function of rainfall in the 48 hours prior to sampling.

Table 12. Regression of Bayou Texar station data as a function of rain. Significant parameter estimates ( $p < 0.06$ ) are indicated in bold.

| station code | station name                          | R <sup>2</sup> | Slope        | p value       | Intercept    | P value         | geomean at 0 rain |
|--------------|---------------------------------------|----------------|--------------|---------------|--------------|-----------------|-------------------|
| 1            | <b>Cervantes Bridge</b>               | <b>0.103</b>   | <b>0.355</b> | <b>0.043</b>  | <b>1.087</b> | <b>7.61E-15</b> | <b>12.213</b>     |
| 2            | Brainerd St. pond                     | 0.021          | 0.345        | 0.205         | <b>1.782</b> | <b>1.08E-20</b> | <b>60.593</b>     |
| 3            | Bayview Park pvc/storm drain          | 0.030          | 0.171        | 0.164         | <b>0.966</b> | <b>8.02E-23</b> | <b>9.237</b>      |
| 4            | Tree Roots                            | 0.141          | 0.334        | 0.064         | <b>0.651</b> | <b>2.52E-07</b> | <b>4.476</b>      |
| 5            | <b>Boathouse (point)</b>              | <b>0.175</b>   | <b>0.432</b> | <b>0.008</b>  | <b>0.768</b> | <b>1.35E-11</b> | <b>5.857</b>      |
| 6            | Rocks/Gazebo                          | 0.072          | 0.239        | 0.195         | <b>0.694</b> | <b>1.75E-07</b> | <b>4.940</b>      |
| 7            | Oriental Garden                       | 0.090          | 0.265        | 0.146         | <b>0.700</b> | <b>1.14E-07</b> | <b>5.015</b>      |
| 8            | <b>South Whaley fragmites</b>         | <b>0.169</b>   | <b>0.370</b> | <b>0.037</b>  | <b>0.673</b> | <b>7.02E-08</b> | <b>4.713</b>      |
| 9            | Whaley Ditch storm drain              | 0.030          | 0.205        | 0.283         | <b>1.029</b> | <b>7.74E-13</b> | <b>10.688</b>     |
| 10           | Birnam Woods green SD                 | 0.080          | 0.284        | 0.172         | <b>1.183</b> | <b>9.22E-11</b> | <b>15.228</b>     |
| 11           | Blackshear Ave. SD                    | 0.066          | 0.320        | 0.023         | <b>1.503</b> | <b>2.19E-33</b> | <b>31.821</b>     |
| 12           | <b>Blanford place FW seep</b>         | <b>0.174</b>   | <b>0.502</b> | <b>0.007</b>  | <b>1.594</b> | <b>1.26E-19</b> | <b>39.233</b>     |
| 13           | 34th St. storm drain                  | 0.044          | 0.277        | 0.303         | <b>1.674</b> | <b>1.05E-11</b> | <b>47.245</b>     |
| 14           | Six Cement poles-tan house            | 0.142          | 0.469        | 0.064         | <b>1.601</b> | <b>7.94E-12</b> | <b>39.893</b>     |
| 15           | <b>Carpenter Creek center</b>         | <b>0.238</b>   | <b>0.702</b> | <b>0.013</b>  | <b>1.825</b> | <b>4.45E-12</b> | <b>66.833</b>     |
| 16           | <b>Driftwood 4 SD</b>                 | <b>0.111</b>   | <b>0.474</b> | <b>0.036</b>  | <b>1.873</b> | <b>5.30E-19</b> | <b>74.714</b>     |
| 17           | Texar Woods SD                        | 0.045          | 0.251        | 0.187         | <b>1.460</b> | <b>1.25E-17</b> | <b>28.830</b>     |
| 18           | <b>Seville Dr. (2) SD</b>             | <b>0.182</b>   | <b>0.515</b> | <b>0.006</b>  | <b>1.662</b> | <b>2.86E-20</b> | <b>45.871</b>     |
| 19           | <b>Banguos Court SD</b>               | <b>0.238</b>   | <b>1.014</b> | <b>0.001</b>  | <b>1.466</b> | <b>1.05E-11</b> | <b>29.226</b>     |
| 20           | <b>Bayou Blvd./Perry SD</b>           | <b>0.178</b>   | <b>0.584</b> | <b>0.007</b>  | <b>1.169</b> | <b>1.72E-13</b> | <b>14.747</b>     |
| 21           | <b>12th Ave. bridge</b>               | <b>0.152</b>   | <b>0.530</b> | <b>0.0001</b> | <b>2.294</b> | <b>1.59E-51</b> | <b>196.602</b>    |
| 22           | 9th Ave.                              | 0.034          | 0.174        | 0.218         | <b>2.427</b> | <b>3.81E-33</b> | <b>266.995</b>    |
| 23           | <b>Airport Blvd.</b>                  | <b>0.215</b>   | <b>0.550</b> | <b>0.007</b>  | <b>2.103</b> | <b>2.30E-20</b> | <b>126.772</b>    |
| 24           | Born Court                            | 0.004          | 0.091        | 0.676         | <b>2.447</b> | <b>2.97E-25</b> | <b>280.159</b>    |
| 25           | Boiling Brook                         | 0.046          | 0.270        | 0.163         | <b>2.392</b> | <b>1.75E-26</b> | <b>246.456</b>    |
| 26           | Sears Warehouse                       | 0.182          | 0.679        | 0.069         | <b>1.806</b> | <b>7.12E-08</b> | <b>64.036</b>     |
| 27           | <b>Interstate 10-Historical Dist.</b> | <b>0.105</b>   | <b>0.425</b> | <b>0.047</b>  | <b>2.034</b> | <b>3.55E-20</b> | <b>108.098</b>    |
| 28           | Olive Road                            | 0.019          | 0.197        | 0.463         | <b>1.898</b> | <b>6.86E-14</b> | <b>79.095</b>     |
| 29           | Walton/Davis                          | 0.130          | 0.458        | 0.129         | <b>2.478</b> | <b>4.84E-12</b> | <b>300.913</b>    |




Figure 35. Regression analysis of *Enterococcus* count dependence on rainfall by station in Bayou Texar.









Figure 35, continued. Regression analysis of *Enterococcus* count dependence on rainfall by station in Bayou Texar.









Figure 35, continued. Regression analysis of *Enterococcus* count dependence on rainfall by station in Bayou Texar.









Figure 35, continued. Regression analysis of *Enterococcus* count dependence on rainfall by station in Bayou Texar.








Figure 35, concluded. Regression analysis of *Enterococcus* count dependence on rainfall by station in Bayou Texar.

# Bayou Texar

## *Enterococcus* Rain Dependence




Figure 36. Distribution of slope values for Bayou Texar station rain dependence.

Bayou Texar  
*Enterococcus* CFU/100 ml  
Geomean at Zero Rainfall




Figure 37. Distribution of y-intercept estimates (*Enterococcus* at zero rainfall) for bayou Texar.

### *Site-specific Ground water sampling and Infrared (IR) camera assisted sampling*

The station-specific nature of the fecal contamination observed in the bayous led us to investigate finer scale loading points to the systems. It was hypothesized that if septic tanks were a major contributor, as they appear to be, then ground water sources to the Bayous needed to be examined. This also was seen as a test of the assumptions that drain field effluents were being diffused and filtered by passage through the vadose zone and with distance through setbacks from surface water as prescribed by current regulations.

In Bayou Grande, extreme low tides in the winter of 2001 were targeted for sampling exposed groundwater flows across the intertidal areas into Bayou Grande. These groundwater discharges were visible as streams flowing out of the shoreline across the intertidal zone (Figure 38), plumes of water emerging from invertebrate burrows under hydrostatic pressure, and as less well defined broad seepage areas emerging from the sands and muds of the intertidal zone. For Bayou Grande, sample locations were identified by eye, and by gauging the rate of fill (if any) in shallow (6-8") holes dug in the upper intertidal zone.

Results from the sampling in Bayou Grande revealed significant fecal contamination of these ground water sources (Table 13). Sample processing for this lot did not anticipate the high numbers of Enterococci found, and many results were qualitatively recorded as "too many to count" (TMTC), or a lawn of bacteria (Lawn) resulting from too many cells in the sample to allow individual colonies to be distinguished. A Lawn would represent greater contamination than a TMTC sample. This artificial upper limit on quantification limits options for quantitative analysis, but the qualitative results are dramatic enough to illustrate the point.



Figure 38. In the north arm of Bayou Grande, a ground water stream is pictured leaving the shoreline below the wetland vegetation under the high tide mark and traversing the intertidal zone.

Table 13. *Enterococcus* counts for sampling of groundwater in Bayou Grande. Surface water stations are indicated by the bold font.

| Sta #     | Latitude       | Longitude      | Notes                                  | Enterococcus (CFU/100 mL) |                                |                         |
|-----------|----------------|----------------|--|---------------------------|--------------------------------|-------------------------|
|           |                |                |  | 11 Jan 01                 | 20 Jan 01 Site Notes           | 20 Jan 01               |
| 1         | 30.3827        | 87.2863        | Middle set of pilings                  | 0                         |                                |                         |
| 3         | 30.3831        | 87.2886        | house #309                             | 12                        |                                |                         |
| 4         | 30.3832        | 87.2890        | house #312                             | 44                        |                                |                         |
| 5         | 30.3832        | 87.2892        | house #313                             | 590                       |                                |                         |
| 6         | 30.3833        | 87.2898        | seep                                   | 34                        |                                |                         |
| 7         | 30.3834        | 87.2898        | stream                                 | 133                       | ~50m further up creek          | Lawn <sup>2</sup>       |
| 8         | 30.3837        | 87.2901        | seep                                   | TMTC <sup>1</sup>         |                                | 90                      |
| 9         | 30.3838        | 87.2900        | seep                                   | 23                        |                                |                         |
| <b>10</b> | <b>30.3835</b> | <b>87.2893</b> | <b>DOH #18</b>                         | <b>99</b>                 |                                | <b>Lawn<sup>2</sup></b> |
| 11        | 30.3838        | 87.2886        | seep                                   | 113                       |                                |                         |
| 12        | 30.3844        | 87.2872        | Drain Pipe                             | 269                       |                                | Lawn <sup>2</sup>       |
| <b>13</b> | <b>30.3846</b> | <b>87.2837</b> | <b>DOH #16</b>                         | <b>326</b>                | <b>~20' from water line</b>    | <b>1216<sup>3</sup></b> |
| 14        | 30.3852        | 87.2838        | Stream                                 | 668                       |                                | TMTC <sup>1</sup>       |
| 15        | 30.3851        | 87.2835        | seep                                   | 435                       |                                | TMTC <sup>1</sup>       |
| 15        | 30.3851        | 87.2835        | seep                                   |                           | ~10' from water/behind seawall | 340 <sup>3</sup>        |
| 16        | 30.3846        | 87.2833        | seep                                   | 18                        |                                |                         |
| 17        | 30.3844        | 87.3332        | seep between seawalls, right pine tree | Lawn <sup>2</sup>         |                                | Lawn <sup>2</sup>       |
| 17        | 30.3844        | 87.3332        | seep between seawalls, right pine tree |                           | ~6" from water                 | Lawn <sup>2</sup>       |
| 18        | 30.3836        | 87.2823        | Plastic 18" Drain pipe                 | 398                       | Waterfowl in area              | TMTC <sup>1</sup>       |
| <b>19</b> | <b>30.3833</b> | <b>87.2812</b> | <b>DOH # 15</b>                        | <b>44</b>                 |                                | <b>1568<sup>3</sup></b> |
| 20        | 30.3846        | 87.2801        | Stream                                 | 80                        |                                | 2600 <sup>3</sup>       |
| 21        | 30.3842        | 87.2803        | Between 1st & 2nd Docks                | Lawn <sup>2</sup>         |                                | TMTC <sup>1</sup>       |
| 21        | 30.3842        | 87.2803        | Between 1st & 2nd Docks ground water   |                           |                                | 36                      |
| 22        | 30.3830        | 87.2809        | Broad seep area, Field dup             | 73                        |                                |                         |
| 22        | 30.3830        | 87.2809        | Broad seep area, Field dup             | 76                        |                                |                         |
| <b>23</b> | <b>30.3823</b> | <b>87.2814</b> | <b>DOH # 14</b>                        | <b>11</b>                 |                                | <b>1716<sup>3</sup></b> |

TMTC<sup>1</sup>: too numerous to count.

Lawn<sup>2</sup>: too many bacteria to form distinct colonies on the plate

<sup>3</sup>: result estimated, greater than 60 colonies per plate counted

The use of the infrared (IR) video camera was very effective at identifying the thermal signal from ground water flow into the cooler winter water of Bayou Chico. Thermal plumes could be found leaving the intertidal zone and mixing in with the bayou water. Figure 39 displays visible and IR images of an intertidal beach area experiencing ground water seepage, and an area of higher thermal signature than the surrounding ground water thermal signature. This technology allows the pinpointing of preferential flow paths for ground water entering the bayou when obvious sources as shown in Figure 43 are not apparent.

Sample locations, physico-chemical properties of water samples and the magnitude of *Enterococcus* counts from sampling sites suggested by the use of IR video listed in Table 14. Plots of *Enterococcus* by salinity show high levels in the Bayou itself and generally low numbers in the ground water samples (Figure 40). Three out of 17 ground water samples were equal or greater than 102 colonies per 100 ml (Table 14), but none were high enough to account for the contamination found in the open bayou water. Two of those three samples had elevated salinities of 8.8, 10.4 ppt (Table 14; Figure 40) relative to the third at 4.5 ppt, suggesting that the contamination could have been from bayou water mixing with the ground water on the previous high tide. Chico Bulge #11 had a count of 232 CFU 100ml<sup>-1</sup> (Figure 40) at relatively low salinity (4.5 ppt) suggesting a ground water source for this contamination. This sample tested positive using the molecular test for fecal *Bacteroides*, indicating a relatively fresh contamination source. A drainage stream to the north near the Navy Boulevard Bridge (Rt. 98) (Figure 41) had elevated *Enterococcus* numbers accounting for one of the two lower surface water *Enterococcus* counts, but a vagrant encampment was the likely source of this contamination.

Residences in the vicinity of the Bayou Grande samples were older and likely had septic tank emplacements prior to current restrictions concerning distances to ground and surface waters. In Bayou Chico, residences along the shoreline up gradient hydrologically to Chico Bulge #11 (Pinewood Avenue) are also served by septic tanks. Elevations for the properties from a USGS topographic map indicate a platform at 10 to 15 ft, and a more rapidly sloping shoreline near the bayou from 10 feet to sea level. A total of 7 properties on the east side (Bayou waterfront) of Pinewood Avenue have had septic tank and drain field inspections between 2000 and 2004. All of these inspections reported sandy loam (one sand), and the observed and seasonal water table in the vicinity of the drain fields as undetectable in a 72" auger sample. Drain field locations were all in excess of 100 ft from the Bayou shoreline. One property owner reported installing a 20 ft irrigation well reaching ground water at 18 ft during a drought year. Total residential units on any of the lots along the waterfront did not exceed the regulatory limits of 2500 gallons per day per acre disposal capacity, minimum 100 ft from open water, and minimum 24 inches of vadose zone beneath the drain field (Escambia County Ordinance 99-23). State prescriptions for septic tank placement are less restrictive with 75 feet required between OSTDs and surface waters (State of Florida Administrative Code Chapter 64E-6.005(3)).




Figure 39. Visible light (top) and infrared (bottom) images of the inter tidal beach of Bayou Chico in winter looking south(left) and north (right) towards the arrow in Figure 28. The two images on the left show warmer ground water seepage out of the intertidal sand and plumes extending into the colder bayou water. The Images on the right display a ground water discharge area with elevated temperature relative to the surrounding groundwater seepage indicating a preferential flow path for groundwater entering the bayou (Chico Bulge #10).

Table 14. Data from IR assisted sampling in Bayou Chico.

| Sampling Date | Station ID           | Latitude | Longitude | Temp* | Salinity (ppt) | O2 (mg/L) | Entero (CFU/100mL) | Water source |
|---------------|----------------------|----------|-----------|-------|----------------|-----------|--------------------|--------------|
| 31-Jan-03     | Chico Bulge #4(pier) | 30.40933 | 87.25887  | 13.6  | 13.1           | 9.6       | 680                | surface      |
| 31-Jan-03     | Chico Bulge #9       | 30.40989 | 87.2593   | 13.6  | 14.8           | 9.5       | 780                | surface      |
| 31-Jan-03     | Bulge Jungle         | 30.41083 | 87.25926  | 12.2  | 12.2           | 6.6       | 240                | surface      |
| 31-Jan-03     | Upper WF 1           | 30.4137  | 87.26029  | 14.3  | 11.9           | 4.4       | 170                | surface      |
| 31-Jan-03     | Chico Bulge #1       | 30.40896 | 87.25865  | 13.9  | 2.85           | 3.3       | 26                 | ground       |
| 31-Jan-03     | Chico Bulge #2       | 30.409   | 87.25079  | 14.3  | 8.2            | 2.6       | 14                 | ground       |
| 28-Jan-03     | White Apartments     | 30.40906 | 87.25748  | 13.8  | 2.3            |           | 35                 | ground       |
| 31-Jan-03     | Chico Bulge #3       | 30.40933 | 87.25904  | 13.7  | 7.8            | 6.1       | 26                 | ground       |
| 31-Jan-03     | Chico Bulge #5       | 30.40949 | 87.25913  | 12.3  | 9              | 1.4       | 66                 | ground       |
| 31-Jan-03     | Chico Bulge #7       | 30.40971 | 87.25945  | 15.5  | 0.42           | 5         | 6                  | ground       |
| 31-Jan-03     | Chico Bulge #6       | 30.40974 | 87.25935  | 13.2  | 4.36           | 1.6       | 10                 | ground       |
| 31-Jan-03     | Chico Bulge #10      | 30.40998 | 87.2595   | 16.8  | 0.81           | 4.5       | 2                  | ground       |
| 31-Jan-03     | Chico Bulge #11      | 30.41021 | 87.25947  | 15.7  | 4.5            | 2.6       | 232                | ground       |
| 31-Jan-03     | Chico Bulge #12      | 30.41038 | 87.25945  | 16.3  | 0.565          | 5.2       | 4                  | ground       |
| 31-Jan-03     | Chico Bulge #12.5    | 30.41055 | 87.25933  | 17.2  | 0.96           | 3.2       | 38                 | ground       |
| 28-Jan-03     | Worm Tube(wreck)     | 30.41273 | 87.25855  | 13.5  | 2.89           | 4         | 56                 | ground       |
| 28-Jan-03     | Pier                 | 30.41303 | 87.25845  | 16.4  | 6.03           | 3.6       | 9                  | ground       |
| 31-Jan-03     | Upper WF 4           | 30.41364 | 87.26102  |       |                |           | 12                 | ground       |
| 31-Jan-03     | Upper WF 2           | 30.41374 | 87.26068  | 15.7  | 8.8            | 6.1       | 102                | ground       |
| 31-Jan-03     | Upper WF 5           | 30.41384 | 87.2618   | 15.2  | 1.99           | 6.8       | 20                 | ground       |
| 31-Jan-03     | Upper WF 3           | 30.41386 | 87.26071  | 15    | 10.4           | 3.5       | 144                | ground       |

\*from sampled water exposed to cold air, does not represent an ambient groundwater temperature.




Figure 40. *Enterococcus* counts relative to Salinity for IR assisted point sampling. Circles are ground water samples. Squares are surface water samples of the Bayou.

# Bayou Chico

## *Enterococcus* in Surface and Ground Water




Figure 41. Bayou Chico *Enterococcus* counts at ground water seeps.

## Discussion and Conclusions

Fecal contamination in the urban bayous of Pensacola Florida was highly localized, as revealed by intensive sampling focused on potential loading zones. Chronically impaired stations, as indicated by high geomeans of *Enterococcus* counts, high minimum counts, and high zero rainfall geomeans, were found in low elevation residential areas serviced by septic tank systems. During non-rain periods, stations along the open bayou shoreline and drainages of the Naval Air Station covered by golf course and woods had relatively minor concentrations of *Enterococci*, as did the lower industrialized portion of Bayou Chico and the lower part of Bayou Texar serviced by central sewer. These areas showed contamination that was more episodic and storm water related, especially the lower part of Bayou Texar, which seems to receive a large influx of storm-related fecal material.

Terrestrial wildlife and domestic animal sources would be dominant contributors to the storm water loadings, especially in Bayou Grande with the undeveloped shoreline. Waterfront dog owners should be encouraged to clean up after their animals to limit this source of runoff and infiltration of fecal matter. Waterfowl should be more of a widespread background source, except in the case of feral waterfowl and other birds being actively fed in residential waterfront areas. Release of feral waterfowl should be discouraged, existing birds removed, and feeding discouraged. Not only do feral waterfowl contribute to fecal contamination, but they also increase nutrients and organic loadings leading to eutrophication and associated water quality problems.

Rain effects on fecal concentrations in all three bayous were most apparent for moderate rainfall levels (~ 1.5 inches within 48 hours). In Bayous Grande and Chico, high rainfall resulted in some dilution of counts in those systems and loss of variation among stations. Moderate rainfall effects were highly localized by station, with rain effects damped by chronic loadings at zero rainfall at some stations. Rain effects and storm water contamination were most apparent in the lower parts of Bayous Chico and Texar, where contamination at zero rainfall was low.

Patterns of fecal contamination in Bayou Grande emerge as unique relative to Bayous Chico and Texar, which share overall loading patterns. However, these overall differences and similarities belie underlying patterns that involve more commonality than differences. All three Bayous have significant land area not covered by residential development relying on septic tanks. In Bayou Grande, this region is occupied by the Naval Air Station, which runs nearly the length of the southern shoreline as golf course and undeveloped wooded areas. This arrangement results in fecal concentrations along the salinity gradient from drainages of the residential areas of the northern shoreline. In Bayou Chico, the lower bayou is dominated by industrial land use and has a relatively minor residential component, resulting in the loadings being more restricted to the fresher reaches of the system. In Bayou Texar, the lower bayou is surrounded by residential development, but it is serviced by central sewer, and the chronically loaded stations are nearly all within Carpenter's Creek, representing the major freshwater inflow to that system. Some of this area is serviced by older sewer lines and lift stations, which may be contributing to the problem.

Direct sampling of ground water seepages found substantial contamination of groundwater at very specific locations along a shoreline in Bayou Grande. However, in an area of Bayou Chico, serviced by septic tanks in good soils, vadose zone distances many fold over the 24" drain field distance requirements, and shoreline setbacks well in excess of the 100ft minimum county requirement, counts were substantially lower. The one high sample from this area of Bayou

Chico (sample #11) does raise a concern, but requires further characterization. This sampling effort targeted ground water discharge at the water table/intertidal zone interface. Subtidal groundwater discharge as a route for fecal loading to these systems remains uncharacterized.

Although variance in the datasets precluded any significant correlations between nutrient concentrations, BOD, and fecal contamination, geospatial visual analysis strongly suggests that elevated nitrogen species and BOD are associated with stations also experiencing chronic fecal contamination. Resolving the fecal loading problems will likely ameliorate eutrophication stress as well.

### References Cited

- Bernhard, A.E. and K.G. Field. 2000. A PCR Assay To Discriminate Human and Ruminant Feces on the Basis of Host Differences in *Bacteroides-Prevotella* Genes Encoding 16S rRNA. *Appl. Envir. Microbiol.* 66: 4571-4574.
- Eaton, A.D., L.S. Clesceri, and A.E. Greenberg. 1995. *Standard Methods for the Examination of Water and Wastewater* 19<sup>th</sup> edition. American Public Health Association; American Water Works Association; Water Environment Federation, Washington, D.C.
- Martin, K. J., L. N. Schneider, R. A. Snyder, J. E. Lepo. *in prep.* Detection of *Bacteroides* types characteristic of human or bovine contamination.
- U.S. EPA. 1983. *Methods for Chemical Analysis of Water and Wastes*, EPA/600/4-79/020, March 1983.
- U.S. EPA. 1993. Total nitrate/nitrite TNO3/NO2 EPA Method 353.2 (Revision 2.0): U.S. EPA National Exposure Research Laboratory, August 1993.
- U.S. EPA. 1997. Enterococci (E); EPA Method 1600, Membrane Filter Test Method for Enterococci in Water, EPA-821-R-97-004, May 1997.

### Acknowledgements

This work was supported by a contract from ECHD to UWF for the analysis of fecal contamination in area waterways. The assistance, support, and patience of the ECHD personnel, especially Robert Merritt, Lisa Lavoie, and Mary Beverly is gratefully acknowledged. Thanks also to Jan McCauley, David Polk, and staff at the WRL of UWF for processing of samples. Dr. Kendall Martin, Lori Phillips, Rachel Brooks, and Eric Brewer conducted the *Bacteroides* testing. Kristal Flanders produced the GIS maps. Dr. Ranga Rao, Director of CEDB, provided valuable editorial assistance and advice.